

**Anti-iNOS Picoband Antibody**  
**Catalog # ABO10066****Specification**

---

**Anti-iNOS Picoband Antibody - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | WB                     |
| Primary Accession | <a href="#">P35228</a> |
| Host              | Rabbit                 |
| Reactivity        | Human                  |
| Clonality         | Polyclonal             |
| Format            | Lyophilized            |

**Description**

Rabbit IgG polyclonal antibody for Nitric oxide synthase, inducible(NOS2) detection. Tested with WB in Human.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-iNOS Picoband Antibody - Additional Information**

**Gene ID** 4843

**Other Names**

Nitric oxide synthase, inducible, 1.14.13.39, Hepatocyte NOS, HEP-NOS, Inducible NO synthase, Inducible NOS, iNOS, NOS type II, Peptidyl-cysteine S-nitrosylase NOS2, NOS2, NOS2A

**Calculated MW**

131117 MW KDa

**Application Details**

Western blot, 0.1-0.5 µg/ml, Human<br>

**Tissue Specificity**

Expressed in the liver, retina, bone cells and airway epithelial cells of the lung. Not expressed in the platelets.

**Protein Name**

Nitric oxide synthase, inducible

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg NaN<sub>3</sub>.

**Immunogen**

A synthetic peptide corresponding to a sequence at the C-terminus of human iNOS (1088-1126aa ARDVAHTLKQLVAAKLKLNEEQVEDYFFQLKSQKRYHED), different from the related mouse sequence by five amino acids, and from the related rat sequence by four amino acids.

**Purification**

Immunogen affinity purified.

**Cross Reactivity**

No cross reactivity with other proteins.

**Storage**

**At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.**

**Anti-iNOS Picoband Antibody - Protein Information**

**Name** NOS2 ([HGNC:7873](#))

**Synonyms** NOS2A

**Function**

Produces nitric oxide (NO) which is a messenger molecule with diverse functions throughout the body (PubMed: [7504305](http://www.uniprot.org/citations/7504305), PubMed: [7531687](http://www.uniprot.org/citations/7531687), PubMed: [7544004](http://www.uniprot.org/citations/7544004), PubMed: [7682706](http://www.uniprot.org/citations/7682706)). In macrophages, NO mediates tumoricidal and bactericidal actions. Also has nitrosylase activity and mediates cysteine S-nitrosylation of cytoplasmic target proteins such as PTGS2/COX2 (By similarity). As component of the iNOS-S100A8/9 transnitrosylase complex involved in the selective inflammatory stimulus-dependent S-nitrosylation of GAPDH on 'Cys-247' implicated in regulation of the GAIT complex activity and probably multiple targets including ANXA5, EZR, MSN and VIM (PubMed: [25417112](http://www.uniprot.org/citations/25417112)). Involved in inflammation, enhances the synthesis of pro-inflammatory mediators such as IL6 and IL8 (PubMed: [19688109](http://www.uniprot.org/citations/19688109)).

**Cellular Location**

Cytoplasm, cytosol. Note=Localizes as discrete foci scattered throughout the cytosol and in the presence of SPSB1 and SPSB4, exhibits a more diffuse cytosolic localization.


**Tissue Location**

Expressed in the liver, retina, bone cells and airway epithelial cells of the lung. Not expressed in the platelets Expressed in chondrocytes (PubMed:7504305)

**Anti-iNOS Picoband Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-iNOS Picoband Antibody - Images**



Western blot analysis of iNOS expression in HELA whole cell lysates (lane 1). iNOS at 130KD was detected using rabbit anti- iNOS Antigen Affinity purified polyclonal antibody (Catalog #ABO10066) at 0.5  $\mu$ g/mL. The blot was developed using chemiluminescence (ECL) method .

#### **Anti-iNOS Picoband Antibody - Background**

Nitric oxide synthase, inducible is an enzyme that in humans is encoded by the NOS2 gene. Nitric oxide (NO) is a messenger molecule with diverse functions throughout the body. In the brain and peripheral nervous system, NO displays many properties of a neurotransmitter; it is implicated in neurotoxicity associated with stroke and neurodegenerative diseases, neural regulation of smooth muscle, including peristalsis, and penile erection. Three different NOS isoforms have been identified which fall into two distinct types, constitutive and inducible. The inducible NOS (iNOS) isoform is expressed in a variety of cell types and tissues in response to inflammatory agents and cytokines. The human iNOS (NOS2) gene is approximately 37 kb in length and consists of 26 exons and 25 introns. NOS2-derived NO is a prerequisite for cytokine signaling and function in innate immunity.