

Anti-MUT Picoband Antibody
Catalog # ABO10146**Specification**

Anti-MUT Picoband Antibody - Product Information

Application	WB, IHC-P
Primary Accession	P22033
Host	Rabbit
Reactivity	Human, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Methylmalonyl-CoA mutase, mitochondrial(MUT) detection.
Tested with WB, IHC-P in Human;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-MUT Picoband Antibody - Additional Information

Gene ID 4594

Other Names

Methylmalonyl-CoA mutase, mitochondrial, MCM, 5.4.99.2, Methylmalonyl-CoA isomerase, MUT

Calculated MW

83134 MW KDa

Application Details

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, By Heat

Western blot, 0.1-0.5 µg/ml, Human, Rat

Subcellular Localization

Mitochondrion matrix.

Protein Name

Methylmalonyl-CoA mutase, mitochondrial

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg NaN₃.

Immunogen

A synthetic peptide corresponding to a sequence at the N-terminus of human MUT (33-65aa LHQQQLHPEWAALAKKQLKGKNPEDLIWHTPE), different from the related mouse sequence by one amino acid.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Anti-MUT Picoband Antibody - Protein Information

Name MMUT ([HGNC:7526](#))

Function

Catalyzes the reversible isomerization of methylmalonyl-CoA (MMCoA) (generated from branched-chain amino acid metabolism and degradation of dietary odd chain fatty acids and cholesterol) to succinyl-CoA (3-carboxypropionyl-CoA), a key intermediate of the tricarboxylic acid cycle.


Cellular Location

Mitochondrion matrix. Mitochondrion. Cytoplasm


Anti-MUT Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-MUT Picoband Antibody - Images


Western blot analysis of MUT expression in rat kidney extract (lane 1) and 22RV1 whole cell lysates (lane 2). MUT at 83KD was detected using rabbit anti- MUT Antigen Affinity purified polyclonal antibody (Catalog #ABO10146) at 0.5 μ g/mL. The blot was developed using chemiluminescence (ECL) method .



MUT was detected in paraffin-embedded sections of human intestinal cancer tissues using rabbit anti- MUT Antigen Affinity purified polyclonal antibody (Catalog # ABO10146) at 1 μ g/mL. The immunohistochemical section was developed using SABC method .



MUT was detected in paraffin-embedded sections of human lung cancer tissues using rabbit anti- MUT Antigen Affinity purified polyclonal antibody (Catalog # ABO10146) at 1 μ g/mL. The immunohistochemical section was developed using SABC method .



MUT was detected in paraffin-embedded sections of human mammary cancer tissues using rabbit anti- MUT Antigen Affinity purified polyclonal antibody (Catalog # ABO10146) at 1 μ g/mL. The immunohistochemical section was developed using SABC method .

Anti-MUT Picoband Antibody - Background

Methylmalonyl-CoA mutase (MUT) is a mitochondrial enzyme that catalyzes the isomerization of methylmalonyl-CoA to succinyl-CoA. This gene is mapped to 6p12.3. MUT requires a vitamin B12-derived prosthetic group, adenosylcobalamin (commonly referred to as AdoCbl), to function. And the product of this enzyme, succinyl-CoA, is a key molecule of the TCA cycle.