

Anti-EPHX2 Picoband Antibody
Catalog # ABO10225**Specification**

Anti-EPHX2 Picoband Antibody - Product Information

Application	WB
Primary Accession	P34913
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Bifunctional epoxide hydrolase 2 (EPHX2) detection. Tested with WB in Human;Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-EPHX2 Picoband Antibody - Additional Information

Gene ID 2053

Other Names

Bifunctional epoxide hydrolase 2, Cytosolic epoxide hydrolase 2, CEH, 3.3.2.10, Epoxide hydratase, Soluble epoxide hydrolase, SEH, Lipid-phosphate phosphatase, 3.1.3.76, EPHX2

Calculated MW

62616 MW KDa

Application Details

Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat

Subcellular Localization

Cytoplasm. Peroxisome.

Protein Name

Bifunctional epoxide hydrolase 2

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Na₃.

Immunogen

A synthetic peptide corresponding to a sequence at the C-terminus of human EPHX2 (505-543aa QHMEDWIPHLKRGHIEDCGHWTQMDKPTEVNQILIKWLD), different from the related mouse sequence by seven amino acids, and from the related rat sequence by eight amino acids.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins.

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Anti-EPHX2 Picoband Antibody - Protein Information

Name EPHX2 ([HGNC:3402](#))

Function

Bifunctional enzyme (PubMed: [12574510](http://www.uniprot.org/citations/12574510)). The C-terminal domain has epoxide hydrolase activity and acts on epoxides (alkene oxides, oxiranes) and arene oxides (PubMed: [12574510](http://www.uniprot.org/citations/12574510), PubMed: [12869654](http://www.uniprot.org/citations/12869654), PubMed: [22798687](http://www.uniprot.org/citations/22798687)). Plays a role in xenobiotic metabolism by degrading potentially toxic epoxides (By similarity). Also determines steady-state levels of physiological mediators (PubMed: [12574510](http://www.uniprot.org/citations/12574510), PubMed: [12869654](http://www.uniprot.org/citations/12869654), PubMed: [21217101](http://www.uniprot.org/citations/21217101), PubMed: [22798687](http://www.uniprot.org/citations/22798687)).


Cellular Location


Cytoplasm. Peroxisome.

Anti-EPHX2 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-EPHX2 Picoband Antibody - Images



Western blot analysis of EPHX2 expression in rat lung extract (lane 1), mouse lung extract (lane 2) and HEPG2 whole cell lysates (lane 3). EPHX2 at 62KD was detected using rabbit anti- EPHX2 Antigen Affinity purified polyclonal antibody (Catalog # ABO10225) at 0.5 μ g/mL. The blot was developed using chemiluminescence (ECL) method .

Anti-EPHX2 Picoband Antibody - Background

Soluble epoxide hydrolase (sEH) is a bifunctional enzyme that in humans is encoded by the EPHX2 gene. It is mapped to 8p21.2-p21.1. This gene encodes a member of the epoxide hydrolase family. The protein, found in both the cytosol and peroxisomes, binds to specific epoxides and converts them to the corresponding dihydrodiols. Mutations in this gene have been associated with familial hypercholesterolemia. Alternatively spliced transcript variants have been described.