

Anti-hnRNP L Picoband Antibody
Catalog # ABO10295**Specification**

Anti-hnRNP L Picoband Antibody - Product Information

Application	WB
Primary Accession	P14866
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Heterogeneous nuclear ribonucleoprotein L (HNRNPL) detection. Tested with WB in Human.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-hnRNP L Picoband Antibody - Additional Information

Gene ID 3191

Other Names

Heterogeneous nuclear ribonucleoprotein L, hnRNP L, HNRNPL, HNRPL

Calculated MW

64133 MW KDa

Application Details

Western blot, 0.1-0.5 µg/ml, Human

Subcellular Localization

Nucleus, nucleoplasm . Cytoplasm . Localized in cytoplasmic mRNP granules containing untranslated mRNAs. These granules are not identical with P bodies or stress granules. .

Protein Name

Heterogeneous nuclear ribonucleoprotein L

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg NaN₃.

Immunogen

A synthetic peptide corresponding to a sequence at the C-terminus of human HnRNP L (546-571aa LLEWESKSDALETGLFLNHYQMKNPN), different from the related mouse and rat sequences by one amino acid.

Purification

Immunogen affinity purified.

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Anti-hnRNP L Picoband Antibody - Protein Information

Name HNRNPL

Synonyms HNRPL

Function

Splicing factor binding to exonic or intronic sites and acting as either an activator or repressor of exon inclusion. Exhibits a binding preference for CA-rich elements (PubMed: [11809897](http://www.uniprot.org/citations/11809897), PubMed: [22570490](http://www.uniprot.org/citations/22570490), PubMed: [24164894](http://www.uniprot.org/citations/24164894), PubMed: [25623890](http://www.uniprot.org/citations/25623890), PubMed: [26051023](http://www.uniprot.org/citations/26051023)). Component of the heterogeneous nuclear ribonucleoprotein (hnRNP) complexes and associated with most nascent transcripts (PubMed: [2687284](http://www.uniprot.org/citations/2687284)). Associates, together with APEX1, to the negative calcium responsive element (nCaRE) B2 of the APEX2 promoter (PubMed: [11809897](http://www.uniprot.org/citations/11809897)). As part of a ribonucleoprotein complex composed at least of ZNF827, HNRNPK and the circular RNA circZNF827 that nucleates the complex on chromatin, may negatively regulate the transcription of genes involved in neuronal differentiation (PubMed: [33174841](http://www.uniprot.org/citations/33174841)). Regulates alternative splicing of a core group of genes involved in neuronal differentiation, likely by mediating H3K36me3-coupled transcription elongation and co-transcriptional RNA processing via interaction with CHD8.


Cellular Location

Nucleus, nucleoplasm. Cytoplasm. Note=Localized in cytoplasmic mRNP granules containing untranslated mRNAs. These granules are not identical with P bodies or stress granules

Anti-hnRNP L Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-hnRNP L Picoband Antibody - Images



Western blot analysis of HnRNP L expression in PANC whole cell lysates (lane 1) and JURKAT whole cell lysates (lane 2). HnRNP L at 64KD was detected using rabbit anti- HnRNP L Antigen Affinity purified polyclonal antibody (Catalog # ABO10295) at 0.5 μ g/mL. The blot was developed using chemiluminescence (ECL) method .

Anti-hnRNP L Picoband Antibody - Background

Heterogeneous nuclear ribonucleoprotein L is a protein that in humans is encoded by the HNRNPL gene. Heterogeneous nuclear RNAs (hnRNAs) which include mRNA precursors and mature mRNAs are associated with specific proteins to form heterogeneous ribonucleoprotein (hnRNP) complexes. HNRNPL is among the proteins that are stably associated with hnRNP complexes and along with other hnRNP proteins is likely to play a major role in the formation, packaging, processing, and function of mRNA. It is present in the nucleoplasm as part of the HNRP complex. HNRP proteins have also been identified outside of the nucleoplasm. Exchange of hnRNP for mRNA-binding proteins accompanies transport of mRNA from the nucleus to the cytoplasm. Since HNRP proteins have been shown to shuttle between the nucleus and the cytoplasm, it is possible that they also have cytoplasmic functions. Two transcript variants encoding different isoforms have been found for this gene.