

**Anti-S100 Beta Antibody**  
**Catalog # ABO10629****Specification****Anti-S100 Beta Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">P04271</a>
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

**Description**

Rabbit IgG polyclonal antibody for Protein S100-B(S100B) detection. Tested with WB in Human;Mouse;Rat.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-S100 Beta Antibody - Additional Information****Gene ID 6285****Other Names**

Protein S100-B, S-100 protein beta chain, S-100 protein subunit beta, S100 calcium-binding protein B, S100B

**Calculated MW**

10713 MW KDa

**Application Details**

Western blot, 0.1-0.5 µg/ml, Human, Rat, Mouse<br>

**Subcellular Localization**

Cytoplasm . Nucleus .

**Tissue Specificity**

Although predominant among the water-soluble brain proteins, S100 is also found in a variety of other tissues.

**Protein Name**

Protein S100-B

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg Thimerosal, 0.05mg NaN3.

**Immunogen**

A synthetic peptide corresponding to a sequence at the C-terminus of human S100 beta(64-78aa DGDGECDFQEFMAFV), identical to the related rat and mouse sequences.

**Purification**

Immunogen affinity purified.

**Cross Reactivity**

No cross reactivity with other proteins

**Storage**

At -20°C for one year. After r° Constitution, at 4°C for one month. It° Can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

**Sequence Similarities**

Belongs to the S-100 family.

**Anti-S100 Beta Antibody - Protein Information**

Name S100B {ECO:0000303|PubMed:6487634, ECO:0000312|HGNC:HGNC:10500}

**Function**

Small zinc- and- and calcium-binding protein that is highly expressed in astrocytes and constitutes one of the most abundant soluble proteins in brain (PubMed:<a href="http://www.uniprot.org/citations/20950652" target="\_blank">20950652</a>, PubMed:<a href="http://www.uniprot.org/citations/6487634" target="\_blank">6487634</a>). Weakly binds calcium but binds zinc very tightly-distinct binding sites with different affinities exist for both ions on each monomer (PubMed:<a href="http://www.uniprot.org/citations/20950652" target="\_blank">20950652</a>, PubMed:<a href="http://www.uniprot.org/citations/6487634" target="\_blank">6487634</a>). Physiological concentrations of potassium ion antagonize the binding of both divalent cations, especially affecting high-affinity calcium-binding sites (By similarity). Acts as a neurotrophic factor that promotes astrogliosis and axonal proliferation (By similarity). Involved in innervation of thermogenic adipose tissue by acting as an adipocyte-derived neurotrophic factor that promotes sympathetic innervation of adipose tissue (By similarity). Binds to and initiates the activation of STK38 by releasing autoinhibitory intramolecular interactions within the kinase (By similarity). Interaction with AGER after myocardial infarction may play a role in myocyte apoptosis by activating ERK1/2 and p53/TP53 signaling (By similarity). Could assist ATAD3A cytoplasmic processing, preventing aggregation and favoring mitochondrial localization (PubMed:<a href="http://www.uniprot.org/citations/20351179" target="\_blank">20351179</a>). May mediate calcium-dependent regulation on many physiological processes by interacting with other proteins, such as TPR-containing proteins, and modulating their activity (PubMed:<a href="http://www.uniprot.org/citations/22399290" target="\_blank">22399290</a>).

**Cellular Location**

Cytoplasm. Nucleus. Secreted {ECO:0000250|UniProtKB:P50114} Note=Secretion into the medium is promoted by interaction with isoform CLSTN3beta of CLSTN3. {ECO:0000250|UniProtKB:P50114}

**Tissue Location**


Although predominant among the water-soluble brain proteins, S100 is also found in a variety of other tissues

**Anti-S100 Beta Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-S100 Beta Antibody - Images



Anti-S100 beta antibody, ABO10629, Western blotting  
Lane 1: Rat Brain Tissue Lysate  
Lane 2: Rat Brain Tissue Lysate  
Lane 3: MCF-7 Cell Lysate  
Lane 4: HELA Cell Lysate  
Lane 5: SMMC Cell Lysate  
Lane 6: JURKAT Cell Lysate  
Lane 7: COLO320 Cell Lysate

### Anti-S100 Beta Antibody - Background

S100 calcium binding protein B or S100B is a protein of the S-100 protein family. S100 genes include at least 13 members which are located as a cluster on chromosome 1q21; however, this gene is located at 21q22.3. S100B is a glial-derived protein that is a well-established biomarker for severity of neurological injury and prognosis for recovery. S100 beta is a calcium-binding protein that is expressed at high levels in brain primarily by astrocytes. Addition of the disulfide-bonded dimeric form of S100 beta to primary neuronal and glial cultures and established cell lines induces axonal extension and alterations in astrocyte proliferation and phenotype, but evidence that S100 beta exerts the same effects *in vivo* has not been presented. Reeves et al.(1994) demonstrated that the same effects of the S100B protein are exerted *in vivo*. They found that both astrogliosis and neurite proliferation occurred in transgenic mice expressing elevated levels of S100b. They suggested that these transgenic mice represent a useful model for studies of the role of S100B in glial-neuronal interactions in normal development and function of the brain and for analyzing the significance of elevated levels of the protein in Down syndrome and Alzheimer disease.