

Anti-Stra8 Antibody
Catalog # ABO10998**Specification**

Anti-Stra8 Antibody - Product Information

Application	WB
Primary Accession	Q7Z7C7
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Stimulated by retinoic acid gene 8 protein homolog(STRA8) detection. Tested with WB in Human.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-Stra8 Antibody - Additional Information

Gene ID 346673

Other Names

Stimulated by retinoic acid gene 8 protein homolog, STRA8 {ECO:0000312|EMBL:AAP47163.1}

Calculated MW

36908 MW KDa

Application Details

Western blot, 0.1-0.5 µg/ml, Human

Subcellular Localization

Cytoplasm .

Tissue Specificity

Expressed specifically in testis. .

Protein Name

Stimulated by retinoic acid gene 8 protein homolog

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg Thimerosal, 0.05mg NaN3.

Immunogen

A synthetic peptide corresponding to a sequence at the C-terminus of human Stra8(313-330aa DEEMIMLQCTETFDDEDL).

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Anti-Stra8 Antibody - Protein Information

Name STRA8 ([HGNC:30653](#))

Function

Meiosis-inducer required for the transition into meiosis for both female and male germ cells. In female germ cells, acts downstream of ZGLP1 as a key effector of the meiotic program: required for premeiotic DNA replication and subsequent events in meiotic prophase. During spermatogenesis, next to its role in meiotic initiation, promotes (but is not required for) spermatogonial differentiation. In complex with MEIOSIN, directly activates the transcription of a subset of critical meiotic genes playing a central role in cell-cycle switching from mitosis to meiosis.

Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:P70278}. Nucleus {ECO:0000250|UniProtKB:P70278}. Note=Shuttles between nucleus and cytoplasm. Nuclear export is XPO1-dependent {ECO:0000250|UniProtKB:P70278}


Tissue Location


Expressed specifically in testis and fetal ovaries.

Anti-Stra8 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Stra8 Antibody - Images



Anti-Stra8 antibody, ABO10998, Western blotting
Lane 1: A231 Cell Lysate
Lane 2: JURKAT Cell Lysate
Lane 3: HT1080 Cell Lysate
Lane 4: SCG Cell Lysate

Anti-Stra8 Antibody - Background

Stra8, stimulated by retinoic acid 8, is expressed in embryonic ovaries just before meiotic initiation, whereas in testis it is expressed after birth. The STRA8 gene contains 9 exons. And this gene is mapped to chromosome 7q31. Retinoic acid(RA) signaling was required for Stra8 expression, and therefore initiation of meiosis, in embryonic mouse ovaries. RA was sufficient to induce Stra8 expression in embryonic testis. Cyp26-mediated RA metabolism was responsible for delaying Stra8 expression in embryonic testis. Since RA signaling regulated Stra8 expression in both embryonic ovaries and adult testis, it was said this portion of the meiotic initiation pathway may be identical in both sexes.