

Anti-AKT3 Antibody
Catalog # ABO11508**Specification**

Anti-AKT3 Antibody - Product Information

Application	WB, IHC-P, ICC
Primary Accession	Q9Y243
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for RAC-gamma serine/threonine-protein kinase(AKT3) detection. Tested with WB, IHC-P, ICC in Human;Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-AKT3 Antibody - Additional Information

Gene ID 10000

Other Names

RAC-gamma serine/threonine-protein kinase, 2.7.11.1, Protein kinase Akt-3, Protein kinase B gamma, PKB gamma, RAC-PK-gamma, STK-2, AKT3, PKBG

Calculated MW

55775 MW KDa

Application Details

Immunocytochemistry , 0.5-1 µg/ml, Mouse, Human, Rat
Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, Rat, Mouse, By Heat
Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat

Subcellular Localization

Nucleus . Cytoplasm . Membrane ; Peripheral membrane protein . Membrane-associated after cell stimulation leading to its translocation.

Tissue Specificity

In adult tissues, it is highly expressed in brain, lung and kidney, but weakly in heart, testis and liver. In fetal tissues, it is highly expressed in heart, liver and brain and not at all in kidney.

Protein Name

RAC-gamma serine/threonine-protein kinase

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Thimerosal, 0.05mg NaN₃.

Immunogen

A synthetic peptide corresponding to a sequence at the N-terminus of human AKT3(122-136aa TSQIDNIGEEEMDAS), identical to the related mouse and rat sequences.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Sequence Similarities

Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family. RAC subfamily.

Anti-AKT3 Antibody - Protein Information

Name AKT3

Synonyms PKBG

Function

AKT3 is one of 3 closely related serine/threonine-protein kinases (AKT1, AKT2 and AKT3) called the AKT kinase, and which regulate many processes including metabolism, proliferation, cell survival, growth and angiogenesis. This is mediated through serine and/or threonine phosphorylation of a range of downstream substrates. Over 100 substrate candidates have been reported so far, but for most of them, no isoform specificity has been reported. AKT3 is the least studied AKT isoform. It plays an important role in brain development and is crucial for the viability of malignant glioma cells. AKT3 isoform may also be the key molecule in up-regulation and down-regulation of MMP13 via IL13. Required for the coordination of mitochondrial biogenesis with growth factor-induced increases in cellular energy demands. Down-regulation by RNA interference reduces the expression of the phosphorylated form of BAD, resulting in the induction of caspase-dependent apoptosis.

Cellular Location

Nucleus. Cytoplasm. Membrane; Peripheral membrane protein Note=Membrane-associated after cell stimulation leading to its translocation

Tissue Location

In adult tissues, it is highly expressed in brain, lung and kidney, but weakly in heart, testis and liver. In fetal tissues, it is highly expressed in heart, liver and brain and not at all in kidney


Anti-AKT3 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)

- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Anti-AKT3 Antibody - Images




Anti-AKT3 antibody, ABO11508, Western blotting
Lane 1: Rat Lung Tissue Lysate
Lane 2: Rat Kidney Tissue Lysate
Lane 3: HELA Cell Lysate
Lane 4: Human Placenta Tissue Lysate
Lane 5: A549 Cell Lysate
Lane 6: NIH3T3 Cell Lysate




Anti-AKT3 antibody, ABO11508, IHC(P)
IHC(P): Rat Kidney Tissue



Anti-AKT3 antibody, ABO11508, IHC(P)IHC(P): Rat Lung Tissue



Anti-AKT3 antibody, ABO11508, ICCICC: HEPA Cell



Anti-AKT3 antibody, ABO11508, IHC(P)IHC(P): Human Mammary Cancer Tissue

Anti-AKT3 Antibody - Background

RAC-gamma serine/threonine-protein kinase, also known as protein kinase Akt-3, is an enzyme that in humans is encoded by the AKT3 gene. This gene is mapped to 1q43-q44. The protein encoded by this gene is a member of the AKT, also called PKB, serine/threonine protein kinase family. AKT

kinases are known to be regulators of cell signaling in response to insulin and growth factors. They are involved in a wide variety of biological processes including cell proliferation, differentiation, apoptosis, tumorigenesis, as well as glycogen synthesis and glucose uptake. This kinase has been shown to be stimulated by platelet-derived growth factor(PDGF), insulin, and insulin-like growth factor 1(IGF1). AKT3 plays an important role in brain development and is crucial for the viability of malignant glioma cells. AKT3 isoform may also be the key molecule in up-regulation and down-regulation of MMP13 via IL13. This gene is required for the coordination of mitochondrial biogenesis with growth factor-induced increases in cellular energy demands.