

**Anti-MPO Antibody**  
**Catalog # ABO11775****Specification**

---

**Anti-MPO Antibody - Product Information**

Application	WB, IHC-P
Primary Accession	<a href="#">P05164</a>
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

**Description**

Rabbit IgG polyclonal antibody for Myeloperoxidase(MPO) detection. Tested with WB, IHC-P in Human;Mouse;Rat.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-MPO Antibody - Additional Information**

**Gene ID** 4353

**Other Names**

Myeloperoxidase, MPO, 1.11.2.2, Myeloperoxidase, 89 kDa myeloperoxidase, 84 kDa myeloperoxidase, Myeloperoxidase light chain, Myeloperoxidase heavy chain, MPO

**Calculated MW**

83869 MW KDa

**Application Details**

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, Mouse, Rat, By Heat<br>Western blot, 0.1-0.5 µg/ml, Human<br>

**Subcellular Localization**

Lysosome.

**Protein Name**

Myeloperoxidase

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg Na<sub>3</sub>.

**Immunogen**

E.coli-derived human Myeloperoxidase recombinant protein (Position:S406-S745). Human Myeloperoxidase shares 90% amino acid (aa) sequence identity with mouse Myeloperoxidase.

**Purification**

Immunogen affinity purified.

**Cross Reactivity**

No cross reactivity with other proteins

**Storage**

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

**Sequence Similarities**

Belongs to the peroxidase family. XPO subfamily.

**Anti-MPO Antibody - Protein Information**

**Name** MPO ([HGNC:7218](#))

**Function**

Part of the host defense system of polymorphonuclear leukocytes. It is responsible for microbicidal activity against a wide range of organisms. In the stimulated PMN, MPO catalyzes the production of hypohalous acids, primarily hypochlorous acid in physiologic situations, and other toxic intermediates that greatly enhance PMN microbicidal activity (PubMed: [9922160](http://www.uniprot.org/citations/9922160)). Mediates the proteolytic cleavage of alpha-1-microglobulin to form t-alpha-1-microglobulin, which potently inhibits oxidation of low-density lipoprotein particles and limits vascular damage (PubMed: [25698971](http://www.uniprot.org/citations/25698971)).


**Cellular Location**


Lysosome.

**Anti-MPO Antibody - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


**Anti-MPO Antibody - Images**




Anti-Myeloperoxidase Picoband antibody, ABO11775-1.JPGIHC(P): Human Appendicitis Tissue



Anti-Myeloperoxidase Picoband antibody, ABO11775-2.JPGIHC(P): Mouse Spleen Tissue



Anti-Myeloperoxidase Picoband antibody, ABO11775-3.JPGIHC(P): Rat Spleen Tissue



Anti-Myeloperoxidase Picoband antibody, ABO11775-4.jpgAll lanes: Anti-Myeloperoxidase(ABO11775) at 0.5ug/mlWB: Recombinant Human Myeloperoxidase Protein 0.5ngPredicted bind size: 37KDObserved bind size: 37KD

#### **Anti-MPO Antibody - Background**

Myeloperoxidase (MPO) is a mammalian phagocyte hemoprotein thought to primarily mediate host defense reactions. It is abundantly expressed in neutrophils and secreted during their activation. Myeloperoxidase is part of the host defense system of human polymorphonuclear leukocytes, responsible for microbicidal activity against a wide range of organisms. It is located in the nucleus as well as in the cytoplasm. Intranuclear MPO may help to protect DNA against damage resulting from oxygen radicals produced during myeloid cell maturation and function. The standard product used in this kit is the product of gene recombination, consisting of 697(A49-S745) amino acids with the molecular mass of 80KDa.