

Anti-Ubiquitin Picoband Antibody
Catalog # ABO11824**Specification**

Anti-Ubiquitin Picoband Antibody - Product Information

Application	WB, IHC-P
Primary Accession	P0CG47
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Ubiquitin-60S ribosomal protein L40(UBA52) detection. Tested with WB, IHC-P in Human;Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-Ubiquitin Picoband Antibody - Additional Information

Gene ID 7314

Other Names

Polyubiquitin-B, Ubiquitin, UBB

Calculated MW

25762 MW KDa

Application Details

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, Mouse, Rat, By Heat
Western blot, 0.1-0.5 µg/ml, Human

Subcellular Localization

Ubiquitin: Cytoplasm . Nucleus .

Protein Name

Polyubiquitin-B

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg NaN₃.

Immunogen

E.coli-derived human Ubiquitin recombinant protein (Position: M77-G152).

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Sequence Similarities

Belongs to the ubiquitin family.

Anti-Ubiquitin Picoband Antibody - Protein Information**Name** UBB**Function**

[Ubiquitin]: Exists either covalently attached to another protein, or free (unanchored). When covalently bound, it is conjugated to target proteins via an isopeptide bond either as a monomer (monoubiquitin), a polymer linked via different Lys residues of the ubiquitin (polyubiquitin chains) or a linear polymer linked via the initiator Met of the ubiquitin (linear polyubiquitin chains). Polyubiquitin chains, when attached to a target protein, have different functions depending on the Lys residue of the ubiquitin that is linked: Lys-6-linked may be involved in DNA repair; Lys-11-linked is involved in ERAD (endoplasmic reticulum-associated degradation) and in cell-cycle regulation; Lys-29-linked is involved in proteotoxic stress response and cell cycle; Lys-33-linked is involved in kinase modification; Lys-48-linked is involved in protein degradation via the proteasome; Lys-63-linked is involved in endocytosis, DNA-damage responses as well as in signaling processes leading to activation of the transcription factor NF-kappa-B. Linear polymer chains formed via attachment by the initiator Met lead to cell signaling. Ubiquitin is usually conjugated to Lys residues of target proteins, however, in rare cases, conjugation to Cys or Ser residues has been observed. When polyubiquitin is free (unanchored-polyubiquitin), it also has distinct roles, such as in activation of protein kinases, and in signaling.


Cellular Location


[Ubiquitin]: Cytoplasm. Nucleus. Mitochondrion outer membrane; Peripheral membrane protein

Anti-Ubiquitin Picoband Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Anti-Ubiquitin Picoband Antibody - Images




Anti- Ubiquitin Picoband antibody, ABO11824, Western blottingAll lanes: Anti Ubiquitin (ABO11824) at 0.5ug/mlWB: Recombinant Human Ubiquitin Protein 0.5ngPredicted bind size: 36KDObserved bind size: 36KD



Anti- Ubiquitin Picoband antibody, ABO11824, IHC(P)IHC(P): Rat Intestine Tissue



Anti- Ubiquitin Picoband antibody, ABO11824, IHC(P)IHC(P): Human Mammary Cancer Tissue



Anti- Ubiquitin Picoband antibody, ABO11824, IHC(P)IHC(P): Mouse Intestine Tissue

Anti-Ubiquitin Picoband Antibody - Background

Ubiquitin (originally, ubiquitous immunopoietic polypeptide) is a small regulatory protein that has been found in almost all tissues (ubiquitously) of eukaryotic organisms. Its gene is mapped to 16p13.3. The ubiquitin protein itself consists of 76 amino acids and has a molecular mass of about 8.5 kDa. Ubiquitin binds to proteins and labels them for destruction. It directs protein recycling and also direct proteins to other locations in the cell, where they control other protein and cell mechanisms.