

**Anti-CD79a Picoband Antibody**  
**Catalog # ABO11863****Specification**

---

**Anti-CD79a Picoband Antibody - Product Information**

Application	WB, IHC-P, ICC
Primary Accession	<a href="#">P11912</a>
Host	Rabbit
Reactivity	Human, Rat
Clonality	Polyclonal
Format	Lyophilized

**Description**

Rabbit IgG polyclonal antibody for B-cell antigen receptor complex-associated protein alpha chain(CD79A) detection. Tested with WB, IHC-P, ICC in Human;Rat.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-CD79a Picoband Antibody - Additional Information**

**Gene ID** 973

**Other Names**

B-cell antigen receptor complex-associated protein alpha chain, Ig-alpha, MB-1 membrane glycoprotein, Membrane-bound immunoglobulin-associated protein, Surface IgM-associated protein, CD79a, CD79A, IGA, MB1

**Calculated MW**

25038 MW KDa

**Application Details**

Immunohistochemistry(Frozen Section), 0.5-1 µg/ml, Human,  
-<br>Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, By  
Heat<br>Western blot, 0.1-0.5 µg/ml, Human, Rat<br>

**Subcellular Localization**

Cell membrane; Single-pass type I membrane protein. Following antigen binding, the BCR has been shown to translocate from detergent-soluble regions of the cell membrane to lipid rafts although signal transduction through the complex can also occur outside lipid rafts. .

**Tissue Specificity**

B-cells.

**Protein Name**

B-cell antigen receptor complex-associated protein alpha chain

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg Na<sub>3</sub>.

**Immunogen**

E.coli-derived human CD79a recombinant protein (Position: T121-P226). Human CD79a shares 91% amino acid (aa) sequence identity with mouse CD79a.

**Purification**

Immunogen affinity purified.

**Cross Reactivity**

No cross reactivity with other proteins

**Storage**

**At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.**

**Sequence Similarities**

Contains 1 Ig-like C2-type (immunoglobulin-like) domain.

**Anti-CD79a Picoband Antibody - Protein Information**

**Name** CD79A

**Synonyms** IGA, MB1

**Function**

Required in cooperation with CD79B for initiation of the signal transduction cascade activated by binding of antigen to the B- cell antigen receptor complex (BCR) which leads to internalization of the complex, trafficking to late endosomes and antigen presentation. Also required for BCR surface expression and for efficient differentiation of pro- and pre-B-cells. Stimulates SYK autophosphorylation and activation. Binds to BLNK, bringing BLNK into proximity with SYK and allowing SYK to phosphorylate BLNK. Also interacts with and increases activity of some Src-family tyrosine kinases. Represses BCR signaling during development of immature B- cells.

**Cellular Location**

Cell membrane; Single-pass type I membrane protein. Note=Following antigen binding, the BCR has been shown to translocate from detergent-soluble regions of the cell membrane to lipid rafts although signal transduction through the complex can also occur outside lipid rafts.

**Tissue Location**


B-cells.

**Anti-CD79a Picoband Antibody - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


## Anti-CD79a Picoband Antibody - Images



Anti- CD79A picoband antibody, ABO11863, Western blottingAll lanes: Anti CD79a (ABO11863) at 0.5ug/mlWB: Recombinant human CD79a Protein 0.5ngPredicted bind size: 45KDObserved bind size: 45KD



Anti- CD79A picoband antibody, ABO11863, Western blottingAll lanes: Anti CD79a (ABO11863) at 0.5ug/mlLane 1: Human Placenta Tissue Lysate at 50ugLane 2: Rat Spleen Tissue Lysate at 50ugPredicted bind size: 25KD Observed bind size: 37KD



Anti- CD79A picoband antibody, ABO11863, IHC(P)IHC(P): Human Tonsil Tissue

#### **Anti-CD79a Picoband Antibody - Background**

Cluster of differentiation CD79A also known as B-cell antigen receptor complex-associated protein alpha chain and MB-1 membrane glycoprotein, is a protein that in humans is encoded by the CD79A gene. It is mapped to 19q13.2. CD79A is a membrane protein with an extracellular immunoglobulin domain, a single span transmembrane region and a short cytoplasmic domain. Genetic deletion of the transmembrane exon of CD79A results in loss of CD79A protein and a complete block of B cell development at the pro to pre B cell transition. Similarly, humans with homozygous splice variants in CD79A predicted to result in loss of the transmembrane region and a truncated or absent protein display agammaglobulinemia and no peripheral B cells.