

Anti-MGMT Picoband Antibody
Catalog # ABO11957**Specification**

Anti-MGMT Picoband Antibody - Product Information

| | |
|-------------------|------------------------|
| Application | WB, IHC-P, ICC |
| Primary Accession | P16455 |
| Host | Rabbit |
| Reactivity | Human |
| Clonality | Polyclonal |
| Format | Lyophilized |

Description

Rabbit IgG polyclonal antibody for Methylated-DNA--protein-cysteine methyltransferase(MGMT) detection. Tested with WB, IHC-P, ICC in Human.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-MGMT Picoband Antibody - Additional Information

Gene ID 4255

Other Names

Methylated-DNA--protein-cysteine methyltransferase, 2.1.1.63, 6-O-methylguanine-DNA methyltransferase, MGMT, O-6-methylguanine-DNA-alkyltransferase, MGMT

Calculated MW

21646 MW KDa

Application Details

Immunocytochemistry , 0.5-1 µg/ml, Human, -
Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, By Heat
Western blot, 0.1-0.5 µg/ml, Human

Subcellular Localization

Nucleus.

Protein Name

Methylated-DNA--protein-cysteine methyltransferase

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Na₃.

Immunogen

E.coli-derived human MGMT recombinant protein (Position: D2-N207). Human MGMT shares 70% and 69% amino acid (aa) sequence identity with mouse and rat MGMT, respectively.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Anti-MGMT Picoband Antibody - Protein Information

Name MGMT

Function

Involved in the cellular defense against the biological effects of O6-methylguanine (O6-MeG) and O4-methylthymine (O4-MeT) in DNA. Repairs the methylated nucleobase in DNA by stoichiometrically transferring the methyl group to a cysteine residue in the enzyme. This is a suicide reaction: the enzyme is irreversibly inactivated.


Cellular Location

Nucleus.

Anti-MGMT Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Anti-MGMT Picoband Antibody - Images

Anti- MGMT Picoband antibody, ABO11957, Western blotting All lanes: Anti MGMT (ABO11957) at 0.5ug/ml WB: Recombinant Human MGMT Protein 0.5ng Predicted bind size: 21KD Observed bind

size: 21KD



Anti- MGMT Picoband antibody, ABO11957, Western blottingAll lanes: Anti MGMT (ABO11957) at 0.5ug/mlLane 1: HELA Whole Cell Lysate at 40ugLane 2: Human Placenta Tissue Lysate at 50ugLane 3: JURKAT Whole Cell Lysate at 40ugPredicted bind size: 21KDObserved bind size: 21KD



Anti- MGMT Picoband antibody, ABO11957, IHC(P)IHC(P): Human Placenta Tissue

Anti-MGMT Picoband Antibody - Background

O6-alkylguanine DNA alkyltransferase (also known as AGT, MGMT or AGAT) is a protein that in humans is encoded by the O6-methylguanine DNA methyltransferase (MGMT) gene. MGMT is mapped to 10q26.3. This gene is crucial for genome stability. It repairs the naturally occurring mutagenic DNA lesion O6-methylguanine back to guanine and prevents mismatch and errors during DNA replication and transcription. It has been found that methylation of the MGMT promoter in the tumor was associated with longer survival.