

Anti-MAOB Picoband Antibody
Catalog # ABO12351**Specification**

Anti-MAOB Picoband Antibody - Product Information

| | |
|-------------------|------------------------|
| Application | WB, IHC-P |
| Primary Accession | P27338 |
| Host | Rabbit |
| Reactivity | Human, Mouse, Rat |
| Clonality | Polyclonal |
| Format | Lyophilized |

Description

Rabbit IgG polyclonal antibody for Amine oxidase [flavin-containing] B(MAOB) detection. Tested with WB, IHC-P in Human;Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-MAOB Picoband Antibody - Additional Information

Gene ID 4129

Other Names

Amine oxidase [flavin-containing] B, 1.4.3.4, Monoamine oxidase type B, MAO-B, MAOB

Calculated MW

58763 MW KDa

Application Details

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, Mouse, Rat, By Heat
Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat

Subcellular Localization

Mitochondrion outer membrane; Single-pass type IV membrane protein; Cytoplasmic side.

Protein Name

Amine oxidase [flavin-containing] B

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Na₃.

Immunogen

A synthetic peptide corresponding to a sequence at the C-terminus of human MAOB (448-484aa REILHAMGKIPEDIWQSEPESVDVPAQPITTTFLER), different from the related mouse sequence by five amino acids, and from the related rat sequence by four amino acids.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Anti-MAOB Picoband Antibody - Protein Information

Name MAOB ([HGNC:6834](#))

Function

Catalyzes the oxidative deamination of primary and some secondary amines such as neurotransmitters, and exogenous amines including the tertiary amine, neurotoxin 1-methyl-4-phenyl-1,2,3,6- tetrahydropyridine (MPTP), with concomitant reduction of oxygen to hydrogen peroxide and participates in the metabolism of neuroactive and vasoactive amines in the central nervous system and peripheral tissues (PubMed:[11049757](http://www.uniprot.org/citations/11049757), PubMed:[11134050](http://www.uniprot.org/citations/11134050), PubMed:[20493079](http://www.uniprot.org/citations/20493079), PubMed:[8316221](http://www.uniprot.org/citations/8316221), PubMed:[8665924](http://www.uniprot.org/citations/8665924)). Preferentially degrades benzylamine and phenylethylamine (PubMed:[11049757](http://www.uniprot.org/citations/11049757), PubMed:[11134050](http://www.uniprot.org/citations/11134050), PubMed:[20493079](http://www.uniprot.org/citations/20493079), PubMed:[8316221](http://www.uniprot.org/citations/8316221), PubMed:[8665924](http://www.uniprot.org/citations/8665924)).


Cellular Location

Mitochondrion outer membrane; Single-pass type IV membrane protein; Cytoplasmic side

Anti-MAOB Picoband Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Anti-MAOB Picoband Antibody - Images




Anti- MAOB Picoband antibody, ABO12351, Western blottingAll lanes: Anti MAOB (ABO12351) at 0.5ug/mlLane 1: Rat Cardiac Muscle Tissue Lysate at 50ugLane 2: Rat Kidney Tissue Lysate at 50ugLane 3: Rat Intestine Tissue Lysate at 50ugLane 4: Mouse Kidney Tissue Lysate at 50ugLane 5: Mouse Intestine Tissue Lysate at 50ugLane 6: Mouse Cardiac Muscle Tissue Lysate at 50ugLane 7: HEPG2 Whole Cell Lysate at 40ugLane 8: HELA Whole Cell Lysate at 40ugLane 9: COLO320 Whole Cell Lysate at 40ugPredicted bind size: 59KDObserved bind size: 59KD




Anti- MAOB Picoband antibody, ABO12351,IHC(P)IHC(P): Mouse Intestine Tissue



Anti- MAOB Picoband antibody, ABO12351,IHC(P)IHC(P): Rat Intestine Tissue



Anti- MAOB Picoband antibody, ABO12351,IHC(P)IHC(P): Human Lung Cancer Tissue



Anti- MAOB Picoband antibody, ABO12351, Western blottingAll lanes: Anti MAOB (ABO12351) at 0.5ug/mlLane 1: Rat Brain Tissue Lysate at 50ugLane 2: Mouse Brain Tissue Lysate at 50ugPredicted bind size: 59KDObserved bind size: 59KD

Anti-MAOB Picoband Antibody - Background

MAOB (MONOAMINE OXIDASE B), also called MAO, BRAIN, AMINE OXIDASE (FLAVIN-CONTAINING) B, is a protein that in humans is encoded by the MAOB gene. MAOB is a member of the flavin monoamine oxidase family. And it is mapped on Xp11.3. MAOB catalyzes the oxidative deamination of biogenic and xenobiotic amines and plays an important role in the metabolism of neuroactive and vasoactive amines in the central nervous system and peripheral tissues. This protein preferentially degrades benzylamine and phenylethylamine. Like MAOA, it also degrades dopamine. MAO-B is involved in the breakdown of dopamine, a neurotransmitter implicated in reinforcing and motivating behaviors as well as movement. MAO-B inhibition is, therefore, associated with enhanced activity of dopamine, as well as with decreased production of hydrogen peroxide, a source of reactive oxygen species.