

**Anti-JNK2 Antibody**  
**Catalog # ABO12755****Specification**

---

**Anti-JNK2 Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">P45984</a>
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized

**Description**

Rabbit IgG polyclonal antibody for Mitogen-activated protein kinase 9(MAPK9) detection. Tested with WB in Human.<br>

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-JNK2 Antibody - Additional Information**

**Gene ID** 5601

**Other Names**

Mitogen-activated protein kinase 9, MAP kinase 9, MAPK 9, 2.7.11.24, JNK-55, Stress-activated protein kinase 1a, SAPK1a, Stress-activated protein kinase JNK2, c-Jun N-terminal kinase 2, MAPK9, JNK2, PRKM9, SAPK1A

**Calculated MW**

48139 MW KDa

**Application Details**

Western blot, 0.1-0.5 µg/ml, Human<br>

**Subcellular Localization**

Cytoplasm . Nucleus .

**Protein Name**

Mitogen-activated protein kinase 9

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg Na<sub>3</sub>.

**Immunogen**

A synthetic peptide corresponding to a sequence in the middle region of human JNK2 (257-288aa RNYVENRPKYPGIKFEELFPDWIFPSESERDK), identical to the related mouse and rat sequences.

**Purification**

Immunogen affinity purified.

**Cross Reactivity**

No cross reactivity with other proteins

**Storage**

**At -20°C for one year. After r<sup>o</sup>Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time.Avoid repeated freezing and thawing.**

**Sequence Similarities**

Belongs to the protein kinase superfamily. CMGC Ser/Thr protein kinase family. MAP kinase subfamily.

**Anti-JNK2 Antibody - Protein Information**

**Name** MAPK9

**Synonyms** JNK2, PRKM9, SAPK1A

**Function**

Serine/threonine-protein kinase involved in various processes such as cell proliferation, differentiation, migration, transformation and programmed cell death (PubMed:<a href="http://www.uniprot.org/citations/10376527" target="\_blank">10376527</a>, PubMed:<a href="http://www.uniprot.org/citations/15805466" target="\_blank">15805466</a>, PubMed:<a href="http://www.uniprot.org/citations/17525747" target="\_blank">17525747</a>, PubMed:<a href="http://www.uniprot.org/citations/19675674" target="\_blank">19675674</a>, PubMed:<a href="http://www.uniprot.org/citations/20595622" target="\_blank">20595622</a>, PubMed:<a href="http://www.uniprot.org/citations/21364637" target="\_blank">21364637</a>, PubMed:<a href="http://www.uniprot.org/citations/22441692" target="\_blank">22441692</a>, PubMed:<a href="http://www.uniprot.org/citations/34048572" target="\_blank">34048572</a>). Extracellular stimuli such as pro- inflammatory cytokines or physical stress stimulate the stress- activated protein kinase/c-Jun N-terminal kinase (SAP/JNK) signaling pathway. In this cascade, two dual specificity kinases MAP2K4/MKK4 and MAP2K7/MKK7 phosphorylate and activate MAPK9/JNK2 (PubMed:<a href="http://www.uniprot.org/citations/10376527" target="\_blank">10376527</a>, PubMed:<a href="http://www.uniprot.org/citations/15805466" target="\_blank">15805466</a>, PubMed:<a href="http://www.uniprot.org/citations/17525747" target="\_blank">17525747</a>, PubMed:<a href="http://www.uniprot.org/citations/19675674" target="\_blank">19675674</a>, PubMed:<a href="http://www.uniprot.org/citations/20595622" target="\_blank">20595622</a>, PubMed:<a href="http://www.uniprot.org/citations/21364637" target="\_blank">21364637</a>, PubMed:<a href="http://www.uniprot.org/citations/22441692" target="\_blank">22441692</a>, PubMed:<a href="http://www.uniprot.org/citations/34048572" target="\_blank">34048572</a>). In turn, MAPK9/JNK2 phosphorylates a number of transcription factors, primarily components of AP-1 such as JUN and ATF2 and thus regulates AP-1 transcriptional activity (PubMed:<a href="http://www.uniprot.org/citations/10376527" target="\_blank">10376527</a>). In response to oxidative or ribotoxic stresses, inhibits rRNA synthesis by phosphorylating and inactivating the RNA polymerase 1-specific transcription initiation factor RRN3 (PubMed:<a href="http://www.uniprot.org/citations/15805466" target="\_blank">15805466</a>). Promotes stressed cell apoptosis by phosphorylating key regulatory factors including TP53 and YAP1 (PubMed:<a href="http://www.uniprot.org/citations/17525747" target="\_blank">17525747</a>, PubMed:<a href="http://www.uniprot.org/citations/21364637" target="\_blank">21364637</a>). In T-cells, MAPK8 and MAPK9 are required for polarized differentiation of T-helper cells into Th1 cells (PubMed:<a href="http://www.uniprot.org/citations/19290929" target="\_blank">19290929</a>). Upon T-cell receptor (TCR) stimulation, is activated by CARMA1, BCL10, MAP2K7 and MAP3K7/TAK1 to regulate JUN protein levels (PubMed:<a href="http://www.uniprot.org/citations/19290929" target="\_blank">19290929</a>). Plays an important role in the osmotic stress- induced epithelial tight-junctions disruption (PubMed:<a

href="http://www.uniprot.org/citations/20595622" target="\_blank">20595622</a>). When activated, promotes beta-catenin/CTNNB1 degradation and inhibits the canonical Wnt signaling pathway (PubMed:<a href="http://www.uniprot.org/citations/19675674" target="\_blank">19675674</a>). Also participates in neurite growth in spiral ganglion neurons (By similarity). Phosphorylates the CLOCK-BMAL1 heterodimer and plays a role in the regulation of the circadian clock (PubMed:<a href="http://www.uniprot.org/citations/22441692" target="\_blank">22441692</a>). Phosphorylates POU5F1, which results in the inhibition of POU5F1's transcriptional activity and enhances its proteasomal degradation (By similarity). Phosphorylates ALKBH5 in response to reactive oxygen species (ROS), promoting ALKBH5 sumoylation and inactivation (PubMed:<a href="http://www.uniprot.org/citations/34048572" target="\_blank">34048572</a>).

#### Cellular Location


Cytoplasm. Nucleus. Note=Colocalizes with POU5F1 in the nucleus.  
{ECO:0000250|UniProtKB:Q9WTU6}

### Anti-JNK2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-JNK2 Antibody - Images



Anti- JNK2 antibody, ABO12755, Western blottingAll lanes: Anti JNK2 (ABO12755) at 0.5ug/mlLane 1: HELA Whole Cell Lysate at 40ugLane 2: COLO320 Whole Cell Lysate at 40ugPredicted bind size: 48KDObserved bind size: 48KD

### Anti-JNK2 Antibody - Background

JNK2 is also known as MAPK9. The protein encoded by this gene is a member of the MAP kinase family. MAP kinases act as an integration point for multiple biochemical signals, and are involved in

a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. This kinase targets specific transcription factors, and thus mediates immediate-early gene expression in response to various cell stimuli. It is most closely related to MAPK8, both of which are involved in UV radiation induced apoptosis, thought to be related to the cytochrome c-mediated cell death pathway. Also, this gene and MAPK8 are also known as c-Jun N-terminal kinases. This kinase blocks the ubiquitination of tumor suppressor p53, and thus it increases the stability of p53 in nonstressed cells. Studies of this gene's mouse counterpart suggest a key role in T-cell differentiation. Several alternatively spliced transcript variants encoding distinct isoforms have been reported.