

Anti-DCX/Doublecortin Rabbit Monoclonal Antibody
Catalog # ABO13271**Specification**

Anti-DCX/Doublecortin Rabbit Monoclonal Antibody - Product Information

Application	WB, FC
Primary Accession	O43602
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

Description

Anti-DCX/Doublecortin Rabbit Monoclonal Antibody . Tested in WB, Flow Cytometry applications.
This antibody reacts with Human, Mouse, Rat.

Anti-DCX/Doublecortin Rabbit Monoclonal Antibody - Additional Information

Gene ID 1641

Other Names

Neuronal migration protein doublecortin, Dublin, Lissencephalin-X, Lis-X, DCX, DBCN, LISX

Calculated MW

40574 MW KDa

Application Details

WB 1:500-1:2000
FC 1:50-1:200

Subcellular Localization

Cytoplasm. Cell projection. Localizes at neurite tips..

Tissue Specificity

Highly expressed in neuronal cells of fetal brain (in the majority of cells of the cortical plate, intermediate zone and ventricular zone), but not expressed in other fetal tissues. In the adult, highly expressed in the brain frontal lobe, but very low expression in other regions of brain, and not detected in heart, placenta, lung, liver, skeletal muscles, kidney and pancreas.

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human DCX

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term

storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-DCX/Doublecortin Rabbit Monoclonal Antibody - Protein Information

Name DCX

Synonyms DBCN, LISX

Function

Microtubule-associated protein required for initial steps of neuronal dispersion and cortex lamination during cerebral cortex development. May act by competing with the putative neuronal protein kinase DCLK1 in binding to a target protein. May in that way participate in a signaling pathway that is crucial for neuronal interaction before and during migration, possibly as part of a calcium ion-dependent signal transduction pathway. May be part with PAFAH1B1/LIS-1 of overlapping, but distinct, signaling pathways that promote neuronal migration.

Cellular Location

Cytoplasm. Cell projection, neuron projection {ECO:0000250|UniProtKB:Q9ESI7}. Note=Localizes at neurite tips. {ECO:0000250|UniProtKB:Q9ESI7}

Tissue Location


Highly expressed in neuronal cells of fetal brain (in the majority of cells of the cortical plate, intermediate zone and ventricular zone), but not expressed in other fetal tissues. In the adult, highly expressed in the brain frontal lobe, but very low expression in other regions of brain, and not detected in heart, placenta, lung, liver, skeletal muscles, kidney and pancreas


Anti-DCX/Doublecortin Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-DCX/Doublecortin Rabbit Monoclonal Antibody - Images





Western blot analysis of DCX expression in C6 cell lysate.