

Anti-CD20 MS4A1 Rabbit Monoclonal Antibody
Catalog # ABO13318**Specification**

Anti-CD20 MS4A1 Rabbit Monoclonal Antibody - Product Information

Application	WB, IHC, FC
Primary Accession	P11836
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid

Description

Anti-CD20 MS4A1 Rabbit Monoclonal Antibody . Tested in WB, IHC, Flow Cytometry applications.
This antibody reacts with Human.

Anti-CD20 MS4A1 Rabbit Monoclonal Antibody - Additional Information

Gene ID 931

Other Names

B-lymphocyte antigen CD20, B-lymphocyte surface antigen B1, Bp35, Leukocyte surface antigen Leu-16, Membrane-spanning 4-domains subfamily A member 1, CD20, MS4A1, CD20

Calculated MW

33077 MW KDa

Application Details

WB 1:500-1:2000
IHC 1:50-1:200
FC 1:50-1:200

Subcellular Localization

Cell membrane ; Multi-pass membrane protein. Cell membrane ; Lipid-anchor.

Tissue Specificity

Expressed on B-cells.

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human CD20

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated

freeze-thaw cycles.

Anti-CD20 MS4A1 Rabbit Monoclonal Antibody - Protein Information

Name MS4A1

Synonyms CD20

Function

B-lymphocyte-specific membrane protein that plays a role in the regulation of cellular calcium influx necessary for the development, differentiation, and activation of B-lymphocytes (PubMed:12920111, PubMed:3925015, PubMed:7684739). Functions as a store-operated calcium (SOC) channel component promoting calcium influx after activation by the B-cell receptor/BCR (PubMed:12920111, PubMed:18474602, PubMed:7684739).

Cellular Location

Cell membrane; Multi-pass membrane protein. Cell membrane; Lipid-anchor. Note=Constitutively associated with membrane rafts.

Tissue Location


Expressed on B-cells.


Anti-CD20 MS4A1 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-CD20 MS4A1 Rabbit Monoclonal Antibody - Images







All lanes use the Antibody at 1:5K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:5K dilution for 1 hour at room temperature.



Western blot analysis of CD20 expression in Raji cell lysate.



Immunohistochemical analysis of paraffin-embedded human tonsil, using CD20 Antibody.