

Anti-Bcl-2 Rabbit Monoclonal Antibody
Catalog # ABO13577**Specification**

Anti-Bcl-2 Rabbit Monoclonal Antibody - Product Information

Application	WB, IHC, IF, ICC, IP, FC
Primary Accession	P10415
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid

Description

Anti-Bcl-2 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications.
This antibody reacts with Human.

Anti-Bcl-2 Rabbit Monoclonal Antibody - Additional Information

Gene ID 596

Other Names

Apoptosis regulator Bcl-2, BCL2

Calculated MW

26266 MW KDa

Application Details

WB 1:500-1:2000
IHC 1:50-1:100
ICC/IF 1:50-1:100
IP 1:50
FC 1:50

Subcellular Localization

Mitochondrion outer membrane ; Single-pass membrane protein. Nucleus membrane ; Single-pass membrane protein. Endoplasmic reticulum membrane ; Single-pass membrane protein.

Tissue Specificity

Expressed in a variety of tissues.

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human Bcl-2

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated

freeze-thaw cycles.

Anti-Bcl-2 Rabbit Monoclonal Antibody - Protein Information

Name BCL2

Function

Suppresses apoptosis in a variety of cell systems including factor-dependent lymphohematopoietic and neural cells (PubMed:1508712, PubMed:8183370). Regulates cell death by controlling the mitochondrial membrane permeability (PubMed:11368354). Appears to function in a feedback loop system with caspases (PubMed:11368354). Inhibits caspase activity either by preventing the release of cytochrome c from the mitochondria and/or by binding to the apoptosis-activating factor (APAF-1) (PubMed:11368354). Also acts as an inhibitor of autophagy: interacts with BECN1 and AMBRA1 during non-starvation conditions and inhibits their autophagy function (PubMed:18570871, PubMed:20889974, PubMed:21358617). May attenuate inflammation by impairing NLRP1-inflammasome activation, hence CASP1 activation and IL1B release (PubMed:17418785).

Cellular Location

Mitochondrion outer membrane; Single-pass membrane protein. Nucleus membrane; Single-pass membrane protein. Endoplasmic reticulum membrane; Single-pass membrane protein. Cytoplasm {ECO:0000250|UniProtKB:P10417}

Tissue Location


Expressed in a variety of tissues.


Anti-Bcl-2 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Bcl-2 Rabbit Monoclonal Antibody - Images







All lanes use the Antibody at 1:1W dilution for 1 hour at room temperature.




Immunohistochemical analysis of paraffin-embedded Rat liver, using the Antibody at 1:50 dilution.




Immunohistochemical analysis of paraffin-embedded Rat stomach, using the Antibody at 1:50 dilution.




Immunohistochemical analysis of paraffin-embedded Human non-Hodgkin's lymphoma, using the Antibody at 1:50 dilution.




Immunohistochemical analysis of paraffin-embedded Human squamous cell carcinoma , using the Antibody at 1:50 dilution.




Immunohistochemical analysis of paraffin-embedded Mouse skin, using the Antibody at 1:50 dilution.



Western blot analysis of Bcl-2 expression in (1) Jurkat cell lysate;(2) MCF-7 cell lysate.



Immunohistochemical analysis of paraffin-embedded human spleen, using Bcl-2 Antibody.



Immunofluorescent analysis of Jurkat cells, using Bcl-2 Antibody.