

Anti-PRAS40 AKT1S1 Rabbit Monoclonal Antibody
Catalog # ABO13879**Specification**

Anti-PRAS40 AKT1S1 Rabbit Monoclonal Antibody - Product Information

Application	WB, IHC, IP, FC
Primary Accession	Q96B36
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human, Mouse
Clonality	Monoclonal
Format	Liquid

Description

Anti-PRAS40 AKT1S1 Rabbit Monoclonal Antibody . Tested in WB, IHC, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse.

Anti-PRAS40 AKT1S1 Rabbit Monoclonal Antibody - Additional Information

Gene ID 84335

Other Names

Proline-rich AKT1 substrate 1, 40 kDa proline-rich AKT substrate, AKT1S1
{ECO:0000312|EMBL:AAH16043.1}

Calculated MW

27383 MW KDa

Application Details

WB 1:500-1:2000
IHC 1:50-1:200
IP 1:50
FC 1:50

Subcellular Localization

Cytoplasm, cytosol. Found in the cytosolic fraction of the brain..

Tissue Specificity

Widely expressed with highest levels of expression in liver and heart. Expressed at higher levels in cancer cell lines (e.g. A-549 and HeLa) than in normal cell lines (e.g. HEK293)..

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human PRAS40

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for

**up to one month. Avoid repeated
freeze-thaw cycles.**

Anti-PRAS40 AKT1S1 Rabbit Monoclonal Antibody - Protein Information

Name AKT1S1 {ECO:0000312|EMBL:AAH16043.1}

Function

Negative regulator of the mechanistic target of rapamycin complex 1 (mTORC1), an evolutionarily conserved central nutrient sensor that stimulates anabolic reactions and macromolecule biosynthesis to promote cellular biomass generation and growth (PubMed:17277771, PubMed:17386266, PubMed:17510057, PubMed:29236692). In absence of insulin and nutrients, AKT1S1 associates with the mTORC1 complex and directly inhibits mTORC1 activity by blocking the MTOR substrate- recruitment site (PubMed:29236692). In response to insulin and nutrients, AKT1S1 dissociates from mTORC1 (PubMed:17386266, PubMed:18372248). Its activity is dependent on its phosphorylation state and binding to 14-3-3 (PubMed:16174443, PubMed:18372248). May also play a role in nerve growth factor-mediated neuroprotection (By similarity).

Cellular Location

Cytoplasm, cytosol {ECO:0000250|UniProtKB:Q9D1F4}. Note=Found in the cytosolic fraction of the brain. {ECO:0000250|UniProtKB:Q9D1F4}

Tissue Location


Widely expressed with highest levels of expression in liver and heart. Expressed at higher levels in cancer cell lines (e.g. A-549 and HeLa) than in normal cell lines (e.g. HEK293)


Anti-PRAS40 AKT1S1 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-PRAS40 AKT1S1 Rabbit Monoclonal Antibody - Images





Western blot analysis of PRAS40 expression in 293T cell lysate.