

Anti-Sonic Hedgehog SHH Rabbit Monoclonal Antibody
Catalog # ABO14176**Specification****Anti-Sonic Hedgehog SHH Rabbit Monoclonal Antibody - Product Information**

Application	WB, IHC, IF, ICC, FC
Primary Accession	Q15465
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid

Description

Anti-Sonic Hedgehog SHH Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human.

Anti-Sonic Hedgehog SHH Rabbit Monoclonal Antibody - Additional Information

Gene ID 6469

Other Names

Sonic hedgehog protein, SHH, 3.1.-., HHG-1, Shh unprocessed N-terminal signaling and C-terminal autoprocessing domains, ShhNC, Sonic hedgehog protein N-product, ShhN, Shh N-terminal processed signaling domains, ShhNp, SHH (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=10848)
HGNC:10848

Calculated MW

49607 MW KDa

Application Details

WB 1:500-1:2000
IHC 1:50-1:200
ICC/IF 1:50-1:200
FC 1:50

Subcellular Localization

Sonic hedgehog protein C-product: Secreted, extracellular space. The C-terminal peptide diffuses from the cell..

Tissue Specificity

Expressed in fetal intestine, liver, lung, and kidney. Not expressed in adult tissues.

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human Sonic Hedgehog

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-Sonic Hedgehog SHH Rabbit Monoclonal Antibody - Protein Information

Name SHH ([HGNC:10848](#))

Function

[Sonic hedgehog protein]: The C-terminal part of the sonic hedgehog protein precursor displays an autoproteolysis and a cholesterol transferase activity (By similarity). Both activities result in the cleavage of the full-length protein into two parts (ShhN and ShhC) followed by the covalent attachment of a cholesterol moiety to the C-terminal of the newly generated ShhN (By similarity). Both activities occur in the endoplasmic reticulum (By similarity). Once cleaved, ShhC is degraded in the endoplasmic reticulum (By similarity).

Cellular Location


[Sonic hedgehog protein]: Endoplasmic reticulum membrane. Golgi apparatus membrane. Secreted Note=Co-localizes with HHAT in the ER and Golgi membrane

Anti-Sonic Hedgehog SHH Rabbit Monoclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Anti-Sonic Hedgehog SHH Rabbit Monoclonal Antibody - Images




All lanes use the Antibody at 1:3K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:3K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:3K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:3K dilution for 1 hour at room temperature.



Western blot analysis of Sonic Hedgehog in HepG2 cell lysate.



Immunohistochemical analysis of paraffin-embedded human endometrium, using Sonic Hedgehog Antibody.