

Anti-Histone H4 (tri methyl K20) HIST1H4A Rabbit Monoclonal Antibody
Catalog # ABO14314**Specification**

Anti-Histone H4 (tri methyl K20) HIST1H4A Rabbit Monoclonal Antibody - Product Information

Application	WB, IHC
Primary Accession	P62805
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

Description

Anti-Histone H4 (tri methyl K20) HIST1H4A Rabbit Monoclonal Antibody . Tested in WB, IHC applications. This antibody reacts with Human, Mouse, Rat.

Anti-Histone H4 (tri methyl K20) HIST1H4A Rabbit Monoclonal Antibody - Additional Information

Gene ID 121504;554313;8294;8359;8360;8361;8362;8363;8364;8365;8366;8367;8368;8370

Other Names

Histone H4, H4C1, H4/A, H4FA, HIST1H4A

Calculated MW

11367 MW KDa

Application Details

WB 1:500-1:2000
IHC 1:50-1:200

Subcellular Localization

Nucleus. Chromosome.

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human Histone H4 (tri methyl K20)

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-Histone H4 (tri methyl K20) HIST1H4A Rabbit Monoclonal Antibody - Protein Information

Name H4C1

Synonyms H4/A, H4FA, HIST1H4A

Function

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

Cellular Location


Nucleus {ECO:0000250|UniProtKB:P62806}. Chromosome. Note=Localized to the nucleus when acetylated in step 11 spermatids. {ECO:0000250|UniProtKB:P62806}

Anti-Histone H4 (tri methyl K20) HIST1H4A Rabbit Monoclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Histone H4 (tri methyl K20) HIST1H4A Rabbit Monoclonal Antibody - Images



Western blot analysis of Histone H4 (tri methyl K20) expression in HeLa cell lysate.



Immunohistochemical analysis of paraffin-embedded human stomach, using Histone H4 (tri methyl K20) Antibody.