

**Anti-SND1 Antibody Picoband™ (monoclonal, 5F5E9)**  
**Catalog # ABO15112****Specification****Anti-SND1 Antibody Picoband™ (monoclonal, 5F5E9) - Product Information**

Application	WB, IHC, IF, ICC, FC
Primary Accession	<a href="#">Q7KZF4</a>
Host	Mouse
Isotype	Mouse IgG2b
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Lyophilized

**Description**

Anti-SND1 Antibody Picoband™ (monoclonal, 5F5E9) . Tested in Flow Cytometry, IF, IHC, ICC, WB applications. This antibody reacts with Human, Mouse, Rat.

**Reconstitution**

Adding 0.2 ml of distilled water will yield a concentration of 500 µg/ml.

**Anti-SND1 Antibody Picoband™ (monoclonal, 5F5E9) - Additional Information**

**Gene ID** 27044

**Other Names**

Staphylococcal nuclease domain-containing protein 1, 3.1.31.1, 100 kDa coactivator, EBNA2 coactivator p100, Tudor domain-containing protein 11, p100 co-activator, SND1, TDRD11

**Calculated MW**

110 kDa KDa

**Application Details**

Western blot, 0.25-0.5 µg/ml, Human, Mouse, Rat<br> Immunohistochemistry(Paraffin-embedded Section), 2-5 µg/ml, Human, Mouse, Rat<br> Immunocytochemistry/Immunofluorescence, 5 µg/ml, Human<br> Flow Cytometry, 1-3 µg/1x10<sup>6</sup> cells, Human<br>

**Contents**

Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na<sub>2</sub>HPO<sub>4</sub>.

**Immunogen**

E.coli-derived human SND1 recombinant protein (Position: Q20-D204).

**Purification**

Immunogen affinity purified.

**Storage**

**At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.**

**Anti-SND1 Antibody Picoband™ (monoclonal, 5F5E9) - Protein Information****Name** SND1**Synonyms** TDRD11**Function**

Endonuclease that mediates miRNA decay of both protein-free and AGO2-loaded miRNAs (PubMed:<a href="http://www.uniprot.org/citations/18453631" target="\_blank">18453631</a>, PubMed:<a href="http://www.uniprot.org/citations/28546213" target="\_blank">28546213</a>). As part of its function in miRNA decay, regulates mRNAs involved in G1-to-S phase transition (PubMed:<a href="http://www.uniprot.org/citations/28546213" target="\_blank">28546213</a>). Functions as a bridging factor between STAT6 and the basal transcription factor (PubMed:<a href="http://www.uniprot.org/citations/12234934" target="\_blank">12234934</a>). Plays a role in PIM1 regulation of MYB activity (PubMed:<a href="http://www.uniprot.org/citations/9809063" target="\_blank">9809063</a>). Functions as a transcriptional coactivator for STAT5 (By similarity).

**Cellular Location**

Cytoplasm. Nucleus. Melanosome Note=In IL-4 stimulated cells colocalizes with STAT6 in the nucleus (PubMed:12234934). Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:17081065)


**Tissue Location**

Ubiquitously expressed.

**Anti-SND1 Antibody Picoband™ (monoclonal, 5F5E9) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-SND1 Antibody Picoband™ (monoclonal, 5F5E9) - Images**




Figure 1. Western blot analysis of SND1 using anti-SND1 antibody (M02602-1).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human Hela whole cell lysates,

Lane 2: human Raji whole cell lysates,

Lane 3: human U87 whole cell lysates,

Lane 4: rat stomach tissue lysates,

Lane 5: rat pancreas tissue lysates,

Lane 6: mouse stomach tissue lysates,

Lane 7: mouse pancreas tissue lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-SND1 antigen affinity purified monoclonal antibody (Catalog # M02602-1) at 0.5 µg/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001) with Tanon 5200 system. A specific band was detected for SND1 at approximately 110 kDa. The expected band size for SND1 is at 102 kDa.




Figure 2. IHC analysis of SND1 using anti-SND1 antibody (M02602-1).

SND1 was detected in a paraffin-embedded section of human Hodgkin's lymphoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 µg/ml mouse anti-SND1 Antibody (M02602-1) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the

chromogen.




Figure 3. IHC analysis of SND1 using anti-SND1 antibody (M02602-1).

SND1 was detected in a paraffin-embedded section of human liver cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 µg/ml mouse anti-SND1 Antibody (M02602-1) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.




Figure 4. IHC analysis of SND1 using anti-SND1 antibody (M02602-1).

SND1 was detected in a paraffin-embedded section of human metaplasia of squamous cells of the renal pelvis tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 µg/ml mouse anti-SND1 Antibody (M02602-1) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.






Figure 5. IHC analysis of SND1 using anti-SND1 antibody (M02602-1).

SND1 was detected in a paraffin-embedded section of human tonsil tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2  $\mu$ g/ml mouse anti-SND1 Antibody (M02602-1) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.




Figure 6. IHC analysis of SND1 using anti-SND1 antibody (M02602-1).

SND1 was detected in a paraffin-embedded section of mouse brain tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2  $\mu$ g/ml mouse anti-SND1 Antibody (M02602-1) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.




Figure 7. IHC analysis of SND1 using anti-SND1 antibody (M02602-1).

SND1 was detected in a paraffin-embedded section of rat brain tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2  $\mu$ g/ml mouse anti-SND1 Antibody (M02602-1) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.




Figure 8. IF analysis of SND1 using anti-SND1 antibody (M02602-1).

SND1 was detected in an immunocytochemical section of PC-3 cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5  $\mu$ g/mL mouse anti-SND1 Antibody (M02602-1) overnight at 4°C. DyLight®488 Conjugated Goat Anti-Mouse IgG (BA1126) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.




Figure 9. Flow Cytometry analysis of Hela cells using anti-SND1 antibody (M02602-1). Overlay histogram showing Hela cells stained with M02602-1 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with mouse anti-SND1 Antibody (M02602-1, 1  $\mu\text{g}/1 \times 10^6$  cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-mouse IgG (BA1126, 5-10  $\mu\text{g}/1 \times 10^6$  cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was mouse IgG (1  $\mu\text{g}/1 \times 10^6$ ) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

#### **Anti-SND1 Antibody Picoband™ (monoclonal, 5F5E9) - Background**

Staphylococcal nuclease domain-containing protein 1 also known as 100 kDa coactivator or Tudor domain-containing protein 11 (TDRD11) is a protein that in humans is encoded by the SND1 gene. This gene encodes a transcriptional co-activator that interacts with the acidic domain of Epstein-Barr virus nuclear antigen 2 (EBNA 2), a transcriptional activator that is required for B-lymphocyte transformation. Other transcription factors that interact with this protein are signal transducers and activators of transcription, STATs. This protein is also thought to be essential for normal cell growth. A similar protein in mammals and other organisms is a component of the RNA-induced silencing complex (RISC).