

Anti-FEN1 Rabbit Monoclonal Antibody
Catalog # ABO15172**Specification**

Anti-FEN1 Rabbit Monoclonal Antibody - Product Information

Application	WB, IHC
Primary Accession	P39748
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

Description

Anti-FEN1 Rabbit Monoclonal Antibody . Tested in WB, IHC applications. This antibody reacts with Human, Mouse, Rat.

Anti-FEN1 Rabbit Monoclonal Antibody - Additional Information

Gene ID 2237

Other Names

Flap endonuclease 1 {ECO:0000255|HAMAP-Rule:MF_03140}, FEN-1
{ECO:0000255|HAMAP-Rule:MF_03140}, 3.1.-.- {ECO:0000255|HAMAP-Rule:MF_03140}, DNase IV,
Flap structure-specific endonuclease 1 {ECO:0000255|HAMAP-Rule:MF_03140}, Maturation factor
1, MF1, hFEN-1, FEN1 {ECO:0000255|HAMAP-Rule:MF_03140}, RAD2

Application Details

WB 1:500-1:2000
IHC 1:100-1:500

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human FEN1

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-FEN1 Rabbit Monoclonal Antibody - Protein Information

Name FEN1 {ECO:0000255|HAMAP-Rule:MF_03140}

Synonyms RAD2**Function**

Structure-specific nuclease with 5'-flap endonuclease and 5'- 3' exonuclease activities involved in DNA replication and repair. During DNA replication, cleaves the 5'-overhanging flap structure that is generated by displacement synthesis when DNA polymerase encounters the 5'-end of a downstream Okazaki fragment. It enters the flap from the 5'-end and then tracks to cleave the flap base, leaving a nick for ligation. Also involved in the long patch base excision repair (LP-BER) pathway, by cleaving within the apurinic/apyrimidinic (AP) site- terminated flap. Acts as a genome stabilization factor that prevents flaps from equilibrating into structures that lead to duplications and deletions. Also possesses 5'-3' exonuclease activity on nicked or gapped double-stranded DNA, and exhibits RNase H activity. Also involved in replication and repair of rDNA and in repairing mitochondrial DNA.


Cellular Location

[Isoform 1]: Nucleus, nucleolus. Nucleus, nucleoplasm. Note=Resides mostly in the nucleoli and relocalizes to the nucleoplasm upon DNA damage

Anti-FEN1 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-FEN1 Rabbit Monoclonal Antibody - Images

Western blot analysis of FEN1 expression in (1) HeLa cell lysate; (2) Daudi cell lysate.