

Anti-Prostatic Acid Phosphatase Rabbit Monoclonal Antibody
Catalog # ABO15388**Specification****Anti-Prostatic Acid Phosphatase Rabbit Monoclonal Antibody - Product Information**

Application	WB, IHC, IF, ICC, IP
Primary Accession	P15309
Host	Rabbit
Isotype	IgG
Reactivity	Human, Mouse
Clonality	Monoclonal
Format	Liquid

Description

Anti-Prostatic Acid Phosphatase Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP applications. This antibody reacts with Human, Mouse.

Anti-Prostatic Acid Phosphatase Rabbit Monoclonal Antibody - Additional Information**Gene ID 55****Other Names**

Prostatic acid phosphatase, PAP, 3.1.3.2, 5'-nucleotidase, 5'-NT, 3.1.3.5, Ecto-5'-nucleotidase, Protein tyrosine phosphatase ACP3, 3.1.3.48, Thiamine monophosphatase, TMPase, PAPf39, ACP3 (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=125), ACPP

Calculated MW

50 kDa KDa

Application Details

WB 1:500-1:2000
IHC 1:50-1:200
ICC/IF 1:50-1:200
IP 1:50

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human Prostatic Acid Phosphatase

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-Prostatic Acid Phosphatase Rabbit Monoclonal Antibody - Protein Information

Name ACP3 ([HGNC:125](#))

Synonyms ACPP

Function

A non-specific tyrosine phosphatase that dephosphorylates a diverse number of substrates under acidic conditions (pH 4-6) including alkyl, aryl, and acyl orthophosphate monoesters and phosphorylated proteins (PubMed:10506173, PubMed:15280042, PubMed:20498373, PubMed:9584846). Has lipid phosphatase activity and inactivates lysophosphatidic acid in seminal plasma (PubMed:10506173, PubMed:15280042).

Cellular Location

[Isoform 1]: Secreted

Tissue Location


Highly expressed in the prostate, restricted to glandular and ductal epithelial cells. Also expressed in bladder, kidney, pancreas, lung, cervix, testis and ovary. Weak expression in a subset of pancreatic islet cells, squamous epithelia, the pilosebaceous unit, colonic neuroendocrine cells and skin adnexal structures. Low expression in prostate carcinoma cells and tissues


Anti-Prostatic Acid Phosphatase Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Prostatic Acid Phosphatase Rabbit Monoclonal Antibody - Images





Western blot analysis of Prostatic Acid Phosphatase expression in human prostate cancer lysate.