

Anti-GGT1 Rabbit Monoclonal Antibody
Catalog # ABO15827**Specification**

Anti-GGT1 Rabbit Monoclonal Antibody - Product Information

Application	WB, IHC
Primary Accession	P19440
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid

Description

Anti-GGT1 Rabbit Monoclonal Antibody . Tested in WB, IHC applications. This antibody reacts with Human.

Anti-GGT1 Rabbit Monoclonal Antibody - Additional Information

Gene ID 2678

Other Names

Glutathione hydrolase 1 proenzyme, 3.4.19.13, Gamma-glutamyltransferase 1, Gamma-glutamyltranspeptidase 1, GGT 1, 2.3.2.2, Leukotriene-C4 hydrolase, 3.4.19.14, CD224, Glutathione hydrolase 1 heavy chain, Glutathione hydrolase 1 light chain, GGT1, GGT

Calculated MW

50-60 kDa KDa

Application Details

WB 1:500-1:2000
IHC 1:50-1:200

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human GGT1

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-GGT1 Rabbit Monoclonal Antibody - Protein Information

Name GGT1**Synonyms** GGT**Function**

Cleaves the gamma-glutamyl bond of extracellular glutathione (gamma-Glu-Cys-Gly), glutathione conjugates (such as maresin conjugate (13R)-S-glutathionyl-(14S)-hydroxy-(4Z,7Z,9E,11E,16Z,19Z)-docosahexaenoate, MCTR1) and other gamma-glutamyl compounds (such as leukotriene C4, LTC4) (PubMed:17924658, PubMed:21447318, PubMed:27791009). The metabolism of glutathione by GGT1 releases free glutamate and the dipeptide cysteinyl-glycine, which is hydrolyzed to cysteine and glycine by dipeptidases (PubMed:27791009). In the presence of high concentrations of dipeptides and some amino acids, can also catalyze a transpeptidation reaction, transferring the gamma-glutamyl moiety to an acceptor amino acid to form a new gamma-glutamyl compound (PubMed:17924658, PubMed:21447318, PubMed:7673200, PubMed:7759490, PubMed:8095045, PubMed:8827453). Contributes to cysteine homeostasis, glutathione homeostasis and in the conversion of the leukotriene LTC4 to LTD4.

Cellular Location

Cell membrane; Single-pass type II membrane protein {ECO:0000250|UniProtKB:P07314}


Tissue Location


Detected in fetal and adult kidney and liver, adult pancreas, stomach, intestine, placenta and lung. There are several other tissue-specific forms that arise from alternative promoter usage but that produce the same protein

Anti-GGT1 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-GGT1 Rabbit Monoclonal Antibody - Images



Western blot analysis of GGT1 expression in Human fetal liver lysate.