

Anti-5HT7 Receptor Rabbit Monoclonal Antibody
Catalog # ABO15924**Specification**

Anti-5HT7 Receptor Rabbit Monoclonal Antibody - Product Information

Application	WB
Primary Accession	P34969
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

Description

Anti-5HT7 Receptor Rabbit Monoclonal Antibody . Tested in WB application. This antibody reacts with Human, Mouse, Rat.

Anti-5HT7 Receptor Rabbit Monoclonal Antibody - Additional Information

Gene ID 3363

Other Names

5-hydroxytryptamine receptor 7, 5-HT-7, 5-HT7, 5-HT-X, Serotonin receptor 7, HTR7

Calculated MW

54 kDa KDa

Application Details

WB 1:500-1:2000

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human 5HT7 Receptor

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-5HT7 Receptor Rabbit Monoclonal Antibody - Protein Information

Name HTR7 {ECO:0000303|Ref.3, ECO:0000312|HGNC:HGNC:5302}

Function

G-protein coupled receptor for 5-hydroxytryptamine (serotonin), a biogenic hormone that functions as a neurotransmitter, a hormone and a mitogen (PubMed:35714614, PubMed:8226867). Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of downstream effectors (PubMed:35714614, PubMed:8226867). HTR7 is coupled to G(s) G alpha proteins and mediates activation of adenylate cyclase activity (PubMed:35714614).

Cellular Location

Cell membrane; Multi-pass membrane protein

Tissue Location


[Isoform A]: Predominant isoform in spleen, caudate and hippocampus. [Isoform D]: Minor isoform in terms of expression.

Anti-5HT7 Receptor Rabbit Monoclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Anti-5HT7 Receptor Rabbit Monoclonal Antibody - Images




All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



Western blot analysis of 5HT7 Receptor expression in U-87MG cell lysate.