

Anti-GAA Rabbit Monoclonal Antibody
Catalog # ABO15967**Specification**

Anti-GAA Rabbit Monoclonal Antibody - Product Information

Application	WB
Primary Accession	P10253
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

Description

Anti-GAA Rabbit Monoclonal Antibody . Tested in WB application. This antibody reacts with Human, Mouse, Rat.

Anti-GAA Rabbit Monoclonal Antibody - Additional Information

Gene ID 2548

Other Names

Lysosomal alpha-glucosidase, 3.2.1.20, Acid maltase, Aglucosidase alfa, 76 kDa lysosomal alpha-glucosidase, 70 kDa lysosomal alpha-glucosidase, GAA

Calculated MW

110 kDa, 95 kDa, 76 kDa KDa

Application Details

WB 1:500-1:2000

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human GAA

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-GAA Rabbit Monoclonal Antibody - Protein Information

Name GAA

Function

Essential for the degradation of glycogen in lysosomes (PubMed:14695532, PubMed:18429042, PubMed:1856189, PubMed:7717400). Has highest activity on alpha-1,4-linked glycosidic linkages, but can also hydrolyze alpha-1,6-linked glucans (PubMed:29061980).


Cellular Location

Lysosome. Lysosome membrane


Anti-GAA Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-GAA Rabbit Monoclonal Antibody - Images

All lanes use the Antibody at 1:5K dilution for 1 hour at room temperature.



Western blot analysis of GAA expression in Human fetal liver lysate.