

Anti-TRAF2 Rabbit Monoclonal Antibody
Catalog # ABO16050**Specification**

Anti-TRAF2 Rabbit Monoclonal Antibody - Product Information

Application	WB, IHC, IF, ICC, IP, FC
Primary Accession	Q12933
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid

Description

Anti-TRAF2 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human.

Anti-TRAF2 Rabbit Monoclonal Antibody - Additional Information

Gene ID 7186

Other Names

TNF receptor-associated factor 2, 2.3.2.27, E3 ubiquitin-protein ligase TRAF2, RING-type E3 ubiquitin transferase TRAF2, Tumor necrosis factor type 2 receptor-associated protein 3, TRAF2, TRAP3

Calculated MW

56 kDa KDa

Application Details

WB 1:500-1:2000
IHC 1:50-1:200
ICC/IF 1:50-1:200
IP 1:50
FC 1:50

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human TRAF2

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-TRAF2 Rabbit Monoclonal Antibody - Protein Information

Name TRAF2 {ECO:0000303|PubMed:28489822, ECO:0000312|HGNC:HGNC:12032}

Function

E3 ubiquitin-protein ligase that regulates activation of NF- kappa-B and JNK and plays a central role in the regulation of cell survival and apoptosis (PubMed:10346818, PubMed:11784851, PubMed:12917689, PubMed:15383523, PubMed:18981220, PubMed:19150425, PubMed:19810754, PubMed:19918265, PubMed:19937093, PubMed:20047764, PubMed:20064526, PubMed:20385093, PubMed:20577214, PubMed:22212761). Catalyzes 'Lys-63'-linked ubiquitination of target proteins, such as BIRC3, IKBKE, MLST8, RIPK1 and TICAM1 (PubMed:23453969, PubMed:28489822). Is an essential constituent of several E3 ubiquitin- protein ligase complexes, where it promotes the ubiquitination of target proteins by bringing them into contact with other E3 ubiquitin ligases (PubMed:15383523, PubMed:18981220). Regulates BIRC2 and BIRC3 protein levels by inhibiting their autoubiquitination and subsequent degradation; this does not depend on the TRAF2 RING-type zinc finger domain (PubMed:11907583, PubMed:19506082). Plays a role in mediating activation of NF-kappa-B by EIF2AK2/PKR (PubMed:15121867). In complex with BIRC2 or BIRC3, promotes ubiquitination of IKBKE (PubMed:23453969). Acts as a regulator of mTORC1 and mTORC2 assembly by mediating 'Lys-63'-linked ubiquitination of MLST8, thereby inhibiting formation of the mTORC2 complex, while facilitating assembly of the mTORC1 complex (PubMed:28489822). Required for normal antibody isotype switching from IgM to IgG (By similarity).

Cellular Location


Cytoplasm

Anti-TRAF2 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-TRAF2 Rabbit Monoclonal Antibody - Images



Western blot analysis of TRAF2 expression in Raji cell lysate.