

Anti-BCLG Rabbit Monoclonal Antibody
Catalog # ABO16061**Specification**

Anti-BCLG Rabbit Monoclonal Antibody - Product Information

Application	WB, IHC, IF, ICC, IP, FC
Primary Accession	Q9BZR8
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

Description

Anti-BCLG Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Anti-BCLG Rabbit Monoclonal Antibody - Additional Information

Gene ID 79370

Other Names

Apoptosis facilitator Bcl-2-like protein 14, Bcl2-L-14, Apoptosis regulator Bcl-G, BCL2L14, BCLG

Calculated MW

40 kDa KDa

Application Details

WB 1:500-1:2000
IHC 1:50-1:200
ICC/IF 1:50-1:200
IP 1:50
FC 1:50

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from BCLG

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-BCLG Rabbit Monoclonal Antibody - Protein Information

Name BCL2L14

Synonyms BCLG**Function**

Plays a role in apoptosis.

Cellular Location

Cytoplasm. [Isoform 2]: Endomembrane system. Note=Predominantly localized to cytosolic organelles


Tissue Location

Isoform 1 is widely expressed. Isoform 2 is testis- specific.

Anti-BCLG Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-BCLG Rabbit Monoclonal Antibody - Images

Western blot analysis of BCLG expression in (1) HeLa cell lysate; (2) NIH/3T3 cell lysate; (3) PC12 cell lysate.