

Anti-BIN1 Rabbit Monoclonal Antibody
Catalog # ABO16556**Specification**

Anti-BIN1 Rabbit Monoclonal Antibody - Product Information

Application	WB, IHC, IF, ICC, FC
Primary Accession	O00499
Host	Rabbit
Isotype	IgG
Reactivity	Human, Mouse
Clonality	Monoclonal
Format	Liquid

Description

Anti-BIN1 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry applications.
This antibody reacts with Human, Mouse.

Anti-BIN1 Rabbit Monoclonal Antibody - Additional Information

Gene ID 274

Other Names

Myc box-dependent-interacting protein 1, Amphiphysin II, Amphiphysin-like protein,
Box-dependent myc-interacting protein 1, Bridging integrator 1, BIN1, AMPHL

Calculated MW

65-80 kDa KDa

Application Details

WB 1:500-1:2000
IHC 1:50-1:200
ICC/IF 1:50-1:200
FC 1:50

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50%
glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human BIN1

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-BIN1 Rabbit Monoclonal Antibody - Protein Information

Name BIN1

Synonyms AMPHL

Function

Is a key player in the control of plasma membrane curvature, membrane shaping and membrane remodeling. Required in muscle cells for the formation of T-tubules, tubular invaginations of the plasma membrane that function in depolarization-contraction coupling (PubMed:24755653). Is a negative regulator of endocytosis (By similarity). Is also involved in the regulation of intracellular vesicles sorting, modulation of BACE1 trafficking and the control of amyloid-beta production (PubMed:27179792). In neuronal circuits, endocytosis regulation may influence the internalization of PHF-tau aggregates (By similarity). May be involved in the regulation of MYC activity and the control cell proliferation (PubMed:8782822). Has actin bundling activity and stabilizes actin filaments against depolymerization in vitro (PubMed:28893863).

Cellular Location

[Isoform BIN1]: Nucleus. Cytoplasm Endosome {ECO:0000250|UniProtKB:O08539}. Cell membrane, sarcolemma, T- tubule {ECO:0000250|UniProtKB:O08839}

Tissue Location


Ubiquitous. Highest expression in the brain and muscle (PubMed:9182667). Expressed in oligodendrocytes (PubMed:27488240). Isoform IIA is expressed only in the brain, where it is detected in the gray matter, but not in the white matter (PubMed:27488240). Isoform BIN1 is widely expressed with highest expression in skeletal muscle.


Anti-BIN1 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-BIN1 Rabbit Monoclonal Antibody - Images





Western blot analysis of BIN1 expression in U87-MG cell lysate.