

**Anti-KLHL12 Antibody Picoband™ (monoclonal, 2G11D1)**  
**Catalog # ABO16625****Specification****Anti-KLHL12 Antibody Picoband™ (monoclonal, 2G11D1) - Product Information**

Application	WB, IHC
Primary Accession	<a href="#">Q53G59</a>
Host	Mouse
Isotype	IgG2a
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Lyophilized

**Description**

Anti-KLHL12 Antibody Picoband™ (monoclonal, 2G11D1) . Tested in IHC, WB applications. This antibody reacts with Human, Mouse, Rat.

**Reconstitution**

Adding 0.2 ml of distilled water will yield a concentration of 500 µg/ml.

**Anti-KLHL12 Antibody Picoband™ (monoclonal, 2G11D1) - Additional Information**

**Gene ID** 59349

**Other Names**

Kelch-like protein 12, CUL3-interacting protein 1 {ECO:0000303|Ref.1}, DKIR homolog, hDKIR, KLHL12, C3IP1 {ECO:0000303|Ref.1}

**Calculated MW**

63 kDa KDa

**Application Details**

Western blot, 0.25-0.5 µg/ml, Human, Mouse, Rat<br> Immunohistochemistry(Paraffin-embedded Section), 2-5 µg/ml, Human<br>

**Contents**

Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na2HPO4.

**Immunogen**

E.coli-derived human KLHL12 recombinant protein (Position: R27-I331).

**Purification**

Immunogen affinity purified.

**Storage**

**At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.**

## Anti-KLHL12 Antibody Picoband™ (monoclonal, 2G11D1) - Protein Information

**Name** KLHL12

**Synonyms** C3IP1 {ECO:0000303|Ref.1}

### Function

Substrate-specific adapter of a BCR (BTB-CUL3-RBX1) E3 ubiquitin ligase complex that acts as a negative regulator of Wnt signaling pathway and ER-Golgi transport (PubMed:<a href="http://www.uniprot.org/citations/22358839" target="\_blank">22358839</a>, PubMed:<a href="http://www.uniprot.org/citations/27565346" target="\_blank">27565346</a>). The BCR(KLHL12) complex is involved in ER-Golgi transport by regulating the size of COPII coats, thereby playing a key role in collagen export, which is required for embryonic stem (ES) cells division: BCR(KLHL12) acts by mediating monoubiquitination of SEC31 (SEC31A or SEC31B) (PubMed:<a href="http://www.uniprot.org/citations/22358839" target="\_blank">22358839</a>, PubMed:<a href="http://www.uniprot.org/citations/27565346" target="\_blank">27565346</a>). The BCR(KLHL12) complex is also involved in neural crest specification: in response to cytosolic calcium increase, interacts with the heterodimer formed with PEF1 and PDCD6/ALG-2, leading to bridge together the BCR(KLHL12) complex and SEC31 (SEC31A or SEC31B), promoting monoubiquitination of SEC31 and subsequent collagen export (PubMed:<a href="http://www.uniprot.org/citations/27716508" target="\_blank">27716508</a>). As part of the BCR(KLHL12) complex, also acts as a negative regulator of the Wnt signaling pathway by mediating ubiquitination and subsequent proteolysis of DVL3 (PubMed:<a href="http://www.uniprot.org/citations/16547521" target="\_blank">16547521</a>). The BCR(KLHL12) complex also mediates polyubiquitination of DRD4 and PEF1, without leading to degradation of these proteins (PubMed:<a href="http://www.uniprot.org/citations/18303015" target="\_blank">18303015</a>, PubMed:<a href="http://www.uniprot.org/citations/20100572" target="\_blank">20100572</a>, PubMed:<a href="http://www.uniprot.org/citations/27716508" target="\_blank">27716508</a>).

### Cellular Location

Cytoplasmic vesicle, COPII-coated vesicle

### Tissue Location


Ubiquitously expressed. Highly expressed in testis and at lower levels in the submandibular salivary gland

## Anti-KLHL12 Antibody Picoband™ (monoclonal, 2G11D1) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-KLHL12 Antibody Picoband™ (monoclonal, 2G11D1) - Images






Figure 2. IHC analysis of KLHL12 using anti-KLHL12 antibody (M08568-1).

KLHL12 was detected in a paraffin-embedded section of human liver cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2  $\mu$ g/ml mouse anti-KLHL12 Antibody (M08568-1) overnight at 4°C. Peroxidase Conjugated Goat Anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Mouse IgG Super Vision Assay Kit (Catalog # SV0001) with DAB as the chromogen.




Figure 3. IHC analysis of KLHL12 using anti-KLHL12 antibody (M08568-1).

KLHL12 was detected in a paraffin-embedded section of human thyroid cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2  $\mu$ g/ml mouse anti-KLHL12 Antibody (M08568-1) overnight at 4°C. Peroxidase Conjugated Goat Anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Mouse IgG Super Vision Assay Kit (Catalog # SV0001) with DAB as the chromogen.

#### **Anti-KLHL12 Antibody Picoband™ (monoclonal, 2G11D1) - Background**

Kelch-like protein 12 is a protein that in humans is encoded by the KLHL12 gene. This gene encodes a member of the KLHL (Kelch-like) family of proteins. This protein has been identified as an autoantigen in the autoimmune disease Sjogren's syndrome and as a potential biomarker in primary biliary cirrhosis. This protein may act as a substrate adaptor of the Cullin-3 ubiquitin ligase complex to promote substrate-specific ubiquitylation. Ubiquitylation by this complex has been shown to regulate the Wnt signaling pathway as well as COPII vesicle coat size. A pseudogene has been identified on chromosome 22. Alternative splicing results in multiple transcript variants.