

Anti-ACBD3 Rabbit Monoclonal Antibody

Catalog # ABO16741

Specification

Anti-ACBD3 Rabbit Monoclonal Antibody - Product Information

Application WB
Primary Accession Q9H3P7
Host Rabbit
Isotype Reactivity Human
Clonality Monoclonal
Format Liquid

Description

Anti-ACBD3 Rabbit Monoclonal Antibody . Tested in WB applications. This antibody reacts with

Anti-ACBD3 Rabbit Monoclonal Antibody - Additional Information

Gene ID 64746

Other Names

Golgi resident protein GCP60, Acyl-CoA-binding domain-containing protein 3, Golgi complex-associated protein 1, GOCAP1, Golgi phosphoprotein 1, GOLPH1, PBR- and PKA-associated protein 7, Peripheral benzodiazepine receptor-associated protein PAP7, Golgi resident protein GCP60, N-terminally processed, ACBD3, GCP60, GOCAP1, GOLPH1

Application Details

WB 1:500-1:2000

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human ACBD3

Purification

Affinity-chromatography

Storage Store at -20°C for one year. For short term

storage and frequent use, store at 4°C for

up to one month. Avoid repeated

freeze-thaw cycles.

Anti-ACBD3 Rabbit Monoclonal Antibody - Protein Information

Name ACBD3



Synonyms GCP60, GOCAP1, GOLPH1

Function

Involved in the maintenance of Golgi structure by interacting with giantin, affecting protein transport between the endoplasmic reticulum and Golgi (PubMed:11590181). Involved in hormone-induced steroid biosynthesis in testicular Leydig cells (By similarity). Recruits PI4KB to the Golgi apparatus membrane; enhances the enzyme activity of PI4KB activity via its membrane recruitment thereby increasing the local concentration of the substrate in the vicinity of the kinase (PubMed:27009356).

Cellular Location

Golgi apparatus membrane; Peripheral membrane protein; Cytoplasmic side. Mitochondrion. Note=Also mitochondrial (via its interaction with PBR).

Tissue Location


Ubiquitous, with highest expression in testis and ovary.

Anti-ACBD3 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-ACBD3 Rabbit Monoclonal Antibody - Images



All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.