

Anti-Hexokinase 1 Antibody (6B8-G11-E7)
Mouse Monoclonal Antibody
Catalog # ABV12061**Specification**

Anti-Hexokinase 1 Antibody (6B8-G11-E7) - Product Information

Application	WB
Primary Accession	P19367
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse IgG1

Anti-Hexokinase 1 Antibody (6B8-G11-E7) - Additional Information**Gene ID** 3098**Application & Usage****WB: MCF7, Jurkat, CHO-K1, C2C12, C6, 3T3
and Hela cell lysates****Other Names**

Hexokinase-1, Brain form hexokinase, Hexokinase type I, HK I

Target/Specificity

Hexokinase-1

Antibody Form

Liquid

Appearance

Colorless liquid

Formulation

In PBS (pH 7.4) containing with 0.02% sodium azide and 50% glycerol

Handling

The antibody solution should be gently mixed before use.

Reconstitution & Storage

-20 °C

Background Descriptions**Precautions**

Anti-Hexokinase 1 Antibody (6B8-G11-E7) is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-Hexokinase 1 Antibody (6B8-G11-E7) - Protein Information

Name HK1 ([HGNC:4922](#))

Function

Catalyzes the phosphorylation of various hexoses, such as D- glucose, D-glucosamine, D-fructose, D-mannose and 2-deoxy-D-glucose, to hexose 6-phosphate (D-glucose 6-phosphate, D-glucosamine 6-phosphate, D-fructose 6-phosphate, D-mannose 6-phosphate and 2-deoxy-D-glucose 6- phosphate, respectively) (PubMed:1637300, PubMed:25316723, PubMed:27374331). Does not phosphorylate N-acetyl-D-glucosamine (PubMed:27374331). Mediates the initial step of glycolysis by catalyzing phosphorylation of D-glucose to D-glucose 6-phosphate (By similarity). Involved in innate immunity and inflammation by acting as a pattern recognition receptor for bacterial peptidoglycan (PubMed:27374331). When released in the cytosol, N-acetyl-D-glucosamine component of bacterial peptidoglycan inhibits the hexokinase activity of HK1 and causes its dissociation from mitochondrial outer membrane, thereby activating the NLRP3 inflammasome (PubMed:27374331).

Cellular Location

Mitochondrion outer membrane; Peripheral membrane protein. Cytoplasm, cytosol. Note=The mitochondrial-binding peptide (MBP) region promotes association with the mitochondrial outer membrane (Probable). Dissociates from the mitochondrial outer membrane following inhibition by N-acetyl-D-glucosamine, leading to relocation to the cytosol (PubMed:27374331).

Tissue Location


Isoform 2: Erythrocyte specific (Ref.6). Isoform 3: Testis-specific (PubMed:10978502). Isoform 4: Testis-specific (PubMed:10978502). {ECO:0000269|PubMed:10978502, ECO:0000269|Ref.6}


Anti-Hexokinase 1 Antibody (6B8-G11-E7) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Hexokinase 1 Antibody (6B8-G11-E7) - Images





Western blot detection of Hexokinase 1 in MCF7, JurKat, CHO-K1, C2C12, CS, 3T3 and HeLa cell lysates using Hexokinase 1 mouse mAb

Anti-Hexokinase 1 Antibody (6B8-G11-E7) - Background

Hexokinases phosphorylate glucose to produce glucose-6-phosphate, the first step in most glucose metabolism pathways. This gene encodes a ubiquitous form of hexokinase which localizes to the outer membrane of mitochondria. Mutations in this gene have been associated with hemolytic anemia due to hexokinase deficiency. Alternative splicing of this gene results in five transcript variants which encode different isoforms, some of which are tissue-specific. Each isoform has a distinct N-terminus; the remainder of the protein is identical among all the isoforms. A sixth transcript variant has been described, but due to the presence of several stop codons, it is not thought to encode a protein.