

p57

Mouse Monoclonal antibody(Mab) Catalog # AD80149

Specification

p57 - Product info

Application Primary Accession Reactivity Host Clonality Calculated MW IHC-P P49918 Human Mouse Monoclonal 32177

p57 - Additional info

Gene ID1028Gene NameCDKN1COther NamesCyclin-dependent kinase inhibitor 1C, Cyclin-dependent kinase inhibitor p57, p57Kip2, CDKN1C, KIP2

Dilution IHC-P~~Ready-to-use

Storage Maintain refrigerated at 2-8°C

Precautions

p57 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

p57 - Protein Information

Name CDKN1C


Synonyms KIP2 Function Potent tight-binding inhibitor of several G1 cyclin/CDK complexes (cyclin E-CDK2, cyclin D2-CDK4, and cyclin A-CDK2) and, to lesser extent, of the mitotic cyclin B-CDC2. Negative regulator of cell proliferation. May play a role in maintenance of the nonproliferative state throughout life. Cellular Location Nucleus. **Tissue Location** Expressed in the heart, brain, lung, skeletal muscle, kidney, pancreas and testis. Expressed in the eye. High levels are seen in the placenta while low levels are seen in the liver.




p57 - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture
- p57 Images



Placenta



Immunohistochemical analysis of paraffin-embedded human placenta tissue using AD80149 performed on the Abcarta® FAIP-30 Fully automated IHC platform.Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a Citrate buffer (pH6. 0).Samples were incubated with primary antibody(Ready-to-use) for 15 min at room temperature. AmpSeeTM Detection Systems[Abcepta:AR005] was used as the secondary antibody.