

COX-2
Rabbit Monoclonal antibody(Mab)
Catalog # AD80426**Specification**

COX-2 - Product info

Application	IHC-P
Primary Accession	P35354
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal
Calculated MW	68996

COX-2 - Additional info

Gene ID	5743
Gene Name	PTGS2

Other Names

Prostaglandin G/H synthase 2, 1.14.99.1, Cyclooxygenase-2, COX-2, PHS II, Prostaglandin H2 synthase 2, PGH synthase 2, PGHS-2, Prostaglandin-endoperoxide synthase 2, PTGS2 (HGNC:9605)

Dilution

IHC-P~~Ready-to-use

Storage

Maintain refrigerated at 2-8°C

Precautions

COX-2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

COX-2 - Protein Information

Name PTGS2 ([HGNC:9605](#))

Synonyms
Function

COX2
Converts arachidonate to prostaglandin H2 (PGH2), a committed step in prostanoid synthesis (PubMed:[26859324](#), PubMed:[27226593](#)). Constitutively expressed in some tissues in physiological conditions, such as the endothelium, kidney and brain, and in pathological conditions, such as in cancer. PTGS2 is responsible for production of inflammatory prostaglandins. Up- regulation of PTGS2 is also associated with increased cell

adhesion, phenotypic changes, resistance to apoptosis and tumor angiogenesis. In cancer cells, PTGS2 is a key step in the production of prostaglandin E2 (PGE2), which plays important roles in modulating motility, proliferation and resistance to apoptosis.

Microsome membrane; Peripheral membrane protein. Endoplasmic reticulum membrane; Peripheral membrane protein


Cellular Location

COX-2 - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

COX-2 - Images



Liver



Immunohistochemical analysis of paraffin-embedded human liver tissue using AD80426 performed on the Abcarta® FAIP-30 Fully automated IHC platform. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a EDTA buffer (pH9.0). Samples were incubated with primary antibody (Ready-to-use) for 15 min at room temperature. AmpSee™ Detection Systems [Abcepta:AR005] was used as the secondary

antibody.