

Functional TRAIL-R2 (human) Antibody, mAb(preservative free)
Catalog # ADP0013**Specification**

Functional TRAIL-R2 (human) Antibody, mAb(preservative free) - Product Information

Application	IHC-P, FC, ICC, IP
Primary Accession	O14763
Reactivity	Human
Host	Purified From Concentrated Hybridoma Tissue Culture Supernatant.
Clonality	Monoclonal
Isotype	Mouse IgG1
Gene Source	Human
Application Note	FC, Functional Application, Inhibition (blocks TRAIL-R2 mediated killing if applied in solution), ICC, IHC-P (15 µg/ml), IP, 47878
Calculated MW	IHC-P ~ ~ N/A
Dilution	FC ~ ~ 1:10 ~ 50
	ICC ~ ~ N/A
	IP ~ ~ N/A

Functional TRAIL-R2 (human) Antibody, mAb(preservative free) - Additional Information**Gene ID** 8795**Other Names**

TRAIL Receptor 2; DR5; KILLER; TNFRSF10B; CD262

Target/Specificity

Recognizes human TRAIL-R2. Does not cross-react with human TRAIL-R1, -R3 or -R4.

Format

Liquid. In PBS containing 10% glycerol and 0.02% sodium azide.

Reconstitution & Storage

Stable for at least 1 year after receipt when stored at -20°C.

Precautions

Functional TRAIL-R2 (human) Antibody, mAb(preservative free) is for research use only and not for use in diagnostic or therapeutic procedures.

Functional TRAIL-R2 (human) Antibody, mAb(preservative free) - Protein Information**Name** TNFRSF10B**Synonyms** DR5, KILLER, TRAILR2, TRICK2, ZTNFR9

Function

Receptor for the cytotoxic ligand TNFSF10/TRAIL (PubMed:10549288). The adapter molecule FADD recruits caspase-8 to the activated receptor. The resulting death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation which initiates the subsequent cascade of caspases (aspartate-specific cysteine proteases) mediating apoptosis. Promotes the activation of NF-kappa-B. Essential for ER stress-induced apoptosis.

Cellular Location

Membrane; Single-pass type I membrane protein.


Tissue Location

Widely expressed in adult and fetal tissues; very highly expressed in tumor cell lines such as HeLaS3, K-562, HL-60, SW480, A-549 and G-361; highly expressed in heart, peripheral blood lymphocytes, liver, pancreas, spleen, thymus, prostate, ovary, uterus, placenta, testis, esophagus, stomach and throughout the intestinal tract; not detectable in brain

Functional TRAIL-R2 (human) Antibody, mAb(preservative free) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Functional TRAIL-R2 (human) Antibody, mAb(preservative free) - Images

Immunohistochemical staining of TRAIL-R2 using anti-TRAIL-R2 (human), mAb (HS201) in formalin-fixed and paraffin-embedded (FFPE) human liver tissue (15 µg/ml).

Functional TRAIL-R2 (human) Antibody, mAb(preservative free) - Background

TRAIL-R2 is a receptor for the cytotoxic ligand TRAIL. The adapter molecule FADD recruits caspase-8 to the activated receptor. The resulting death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation which initiates the subsequent cascade of caspases (aspartate-specific cysteine proteases) mediating apoptosis. Promotes the activation of NF-kappa.