

Goat Anti-CD20 / MS4A1 (C Terminus) Antibody
Peptide-affinity purified goat antibody
Catalog # AF1214a**Specification**

Goat Anti-CD20 / MS4A1 (C Terminus) Antibody - Product Information

Application	WB, E
Primary Accession	P11836
Other Accession	NP_690605 , 931
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	33077

Goat Anti-CD20 / MS4A1 (C Terminus) Antibody - Additional Information**Gene ID** 931**Other Names**

B-lymphocyte antigen CD20, B-lymphocyte surface antigen B1, Bp35, Leukocyte surface antigen Leu-16, Membrane-spanning 4-domains subfamily A member 1, CD20, MS4A1, CD20

Dilution

WB~~1:1000

E~~N/A

Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-CD20 / MS4A1 (C Terminus) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-CD20 / MS4A1 (C Terminus) Antibody - Protein Information**Name** MS4A1**Synonyms** CD20**Function**

B-lymphocyte-specific membrane protein that plays a role in the regulation of cellular calcium influx necessary for the development, differentiation, and activation of B-lymphocytes (PubMed:12920111, PubMed:3925015, PubMed:7684739). Functions as a store-operated calcium (SOC) channel component promoting calcium influx after activation by the B-cell receptor/BCR (PubMed:12920111, PubMed:18474602, PubMed:7684739).

Cellular Location

Cell membrane; Multi-pass membrane protein. Cell membrane; Lipid-anchor. Note=Constitutively associated with membrane rafts.

Tissue Location


Expressed on B-cells.

Goat Anti-CD20 / MS4A1 (C Terminus) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat Anti-CD20 / MS4A1 (C Terminus) Antibody - Images



AF1214a (0.05 µg/ml) staining of Human Lymph Node lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Goat Anti-CD20 / MS4A1 (C Terminus) Antibody - Background

This gene encodes a member of the membrane-spanning 4A gene family. Members of this nascent protein family are characterized by common structural features and similar intron/exon splice boundaries and display unique expression patterns among hematopoietic cells and nonlymphoid tissues. This gene encodes a B-lymphocyte surface molecule which plays a role in the development and differentiation of B-cells into plasma cells. This family member is localized to 11q12, among a cluster of family members. Alternative splicing of this gene results in two transcript variants which encode the same protein.

Goat Anti-CD20 / MS4A1 (C Terminus) Antibody - References

Double-Hit mature B-cell lymphomas show a common immunophenotype by flow cytometry that includes decreased CD20 expression. Wu D, et al. Am J Clin Pathol, 2010 Aug. PMID 20660329.

In vivo cytotoxicity of type I CD20 antibodies critically depends on Fc receptor ITAM signaling. de Haij S, et al. Cancer Res, 2010 Apr 15. PMID 20354182.

CD20 as a target for therapeutic type I and II monoclonal antibodies. Beers SA, et al. Semin Hematol, 2010 Apr. PMID 20350657.

New genetic associations detected in a host response study to hepatitis B vaccine. Davila S, et al. Genes Immun, 2010 Apr. PMID 20237496.

CD20 deficiency in humans results in impaired T cell-independent antibody responses. Kuijpers TW, et al. J Clin Invest, 2010 Jan 4. PMID 20038800.