

Goat Anti-ERP29 Antibody
Peptide-affinity purified goat antibody
Catalog # AF1382a**Specification**

Goat Anti-ERP29 Antibody - Product Information

Application	WB, E
Primary Accession	P30040
Other Accession	NP_006808 , 10961 , 67397 (mouse) , 117030 (rat)
Reactivity	Human
Predicted	Rat, Pig, Dog
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	28993

Goat Anti-ERP29 Antibody - Additional Information**Gene ID** 10961**Other Names**

Endoplasmic reticulum resident protein 29, ERp29, Endoplasmic reticulum resident protein 28, ERp28, Endoplasmic reticulum resident protein 31, ERp31, ERP29, C12orf8, ERP28

Dilution

WB~~1:1000

E~~N/A

Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-ERP29 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-ERP29 Antibody - Protein Information**Name** ERP29**Synonyms** C12orf8, ERP28

Function

Does not seem to be a disulfide isomerase. Plays an important role in the processing of secretory proteins within the endoplasmic reticulum (ER), possibly by participating in the folding of proteins in the ER.

Cellular Location

Endoplasmic reticulum lumen. Melanosome. Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV

Tissue Location

Ubiquitous. Mostly expressed in secretory tissues.

Goat Anti-ERP29 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat Anti-ERP29 Antibody - Images

AF1382a (0.1 µg/ml) staining of Human Liver lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Goat Anti-ERP29 Antibody - Background

This gene encodes a reticuloplasmin, a protein which resides in the lumen of the endoplasmic reticulum (ER). The protein shows sequence similarity to the protein disulfide isomerase family. However, it lacks the thioredoxin motif characteristic of this family, suggesting that this protein does not function as a disulfide isomerase. The protein dimerizes and is thought to play a role in the processing of secretory proteins within the ER. Alternative splicing results in multiple transcript variants encoding different isoforms.

Goat Anti-ERP29 Antibody - References

Overexpression of endoplasmic reticulum protein 29 regulates mesenchymal-epithelial transition

and suppresses xenograft tumor growth of invasive breast cancer cells. Bambang IF, et al. Lab Invest, 2009 Nov. PMID 19770839.

ERp29 restricts Connexin43 oligomerization in the endoplasmic reticulum. Das S, et al. Mol Biol Cell, 2009 May. PMID 19321666.

Cytokeratin 19 regulates endoplasmic reticulum stress and inhibits ERp29 expression via p38 MAPK/XBP-1 signaling in breast cancer cells. Bambang IF, et al. Exp Cell Res, 2009 Jul 1. PMID 19265690.

Crystal structure and functional analysis of the protein disulfide isomerase-related protein ERp29. Barak NN, et al. J Mol Biol, 2009 Feb 6. PMID 19084538.

ERp29, an endoplasmic reticulum secretion factor is involved in the growth of breast tumor xenografts. Mkrtchian S, et al. Mol Carcinog, 2008 Nov. PMID 18395818.