

Goat Anti-HIP-55 / SH3P7 Antibody
Peptide-affinity purified goat antibody
Catalog # AF1528a**Specification**

Goat Anti-HIP-55 / SH3P7 Antibody - Product Information

Application	WB, E
Primary Accession	O9UJU6
Other Accession	NP_001116428 , 28988 , 13169 (mouse) , 83527 (rat)
Reactivity	Human, Mouse
Predicted	Rat, Dog
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	48207

Goat Anti-HIP-55 / SH3P7 Antibody - Additional Information**Gene ID** 28988**Other Names**

Drebrin-like protein, Cervical SH3P7, Cervical mucin-associated protein, Drebrin-F, HPK1-interacting protein of 55 kDa, HIP-55, SH3 domain-containing protein 7, DBNL, CMAP, SH3P7

Dilution

WB~~1:1000

E~~N/A

Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-HIP-55 / SH3P7 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-HIP-55 / SH3P7 Antibody - Protein Information**Name** DBNL**Synonyms** CMAP, SH3P7

Function

Adapter protein that binds F-actin and DNM1, and thereby plays a role in receptor-mediated endocytosis. Plays a role in the reorganization of the actin cytoskeleton, formation of cell projections, such as neurites, in neuron morphogenesis and synapse formation via its interaction with WASL and COBL. Does not bind G-actin and promote actin polymerization by itself. Required for the formation of organized podosome rosettes (By similarity). May act as a common effector of antigen receptor-signaling pathways in leukocytes. Acts as a key component of the immunological synapse that regulates T-cell activation by bridging TCRs and the actin cytoskeleton to gene activation and endocytic processes.


Cellular Location


Cytoplasm, cytoskeleton {ECO:0000250|UniProtKB:Q62418}. Cell projection, lamellipodium {ECO:0000250|UniProtKB:Q62418}. Cell projection, ruffle {ECO:0000250|UniProtKB:Q62418}. Cytoplasm, cell cortex {ECO:0000250|UniProtKB:Q62418}. Cytoplasm, cytosol {ECO:0000250|UniProtKB:Q9JHL4}. Synapse {ECO:0000250|UniProtKB:Q62418} Perikaryon {ECO:0000250|UniProtKB:Q62418}. Cell projection, neuron projection {ECO:0000250|UniProtKB:Q62418}. Cell membrane; Peripheral membrane protein {ECO:0000250|UniProtKB:Q62418}; Cytoplasmic side {ECO:0000250|UniProtKB:Q62418}. Cytoplasmic vesicle, clathrin-coated vesicle membrane {ECO:0000250|UniProtKB:Q62418}; Peripheral membrane protein {ECO:0000250|UniProtKB:Q62418}; Cytoplasmic side {ECO:0000250|UniProtKB:Q62418}. Golgi apparatus membrane {ECO:0000250|UniProtKB:Q62418}; Peripheral membrane protein {ECO:0000250|UniProtKB:Q62418}; Cytoplasmic side {ECO:0000250|UniProtKB:Q62418}. Cell projection, podosome {ECO:0000250|UniProtKB:Q62418}. Early endosome. Cell projection, dendrite {ECO:0000250|UniProtKB:Q9JHL4}. Postsynaptic density {ECO:0000250|UniProtKB:Q9JHL4}. Note=Associates with lamellipodial actin and membrane ruffles. Colocalizes with actin and cortactin at podosome dots and podosome rosettes. {ECO:0000250|UniProtKB:Q62418, ECO:0000250|UniProtKB:Q9JHL4}

Goat Anti-HIP-55 / SH3P7 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat Anti-HIP-55 / SH3P7 Antibody - Images



AF1528a staining (1 μ g/ml) of Jurkat lysate (RIPA buffer, 35 μ g total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.

Goat Anti-HIP-55 / SH3P7 Antibody - References

An empirical framework for binary interactome mapping. Venkatesan K, et al. Nat Methods, 2009 Jan. PMID 19060904.

The actin-binding protein Abp1 controls dendritic spine morphology and is important for spine head and synapse formation. Haeckel A, et al. J Neurosci, 2008 Oct 1. PMID 18829961.

Large-scale mapping of human protein-protein interactions by mass spectrometry. Ewing RM, et al. Mol Syst Biol, 2007. PMID 17353931.

Abl-SH3 binding protein 2, 3BP2, interacts with CIN85 and HIP-55. Le Bras S, et al. FEBS Lett, 2007 Mar 6. PMID 17306257.

Global, in vivo, and site-specific phosphorylation dynamics in signaling networks. Olsen JV, et al. Cell, 2006 Nov 3. PMID 17081983.