

**GRB10 Antibody (internal region)**  
**Peptide-affinity purified goat antibody**  
**Catalog # AF2737a****Specification**

---

**GRB10 Antibody (internal region) - Product Information**

Application	WB, E
Primary Accession	<a href="#">Q13322</a>
Other Accession	<a href="#">NP_005302.3</a> , <a href="#">NP_001001549.1</a> , <a href="#">2887</a>
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	67231

**GRB10 Antibody (internal region) - Additional Information****Gene ID** 2887**Other Names**

Growth factor receptor-bound protein 10, GRB10 adapter protein, Insulin receptor-binding protein Grb-IR, GRB10, GRBIR, KIAA0207

**Dilution**

WB~~1:1000

E~~N/A

**Format**

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

GRB10 Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

**GRB10 Antibody (internal region) - Protein Information****Name** GRB10**Synonyms** GRBIR, KIAA0207**Function**

Adapter protein which modulates coupling of a number of cell surface receptor kinases with

specific signaling pathways. Binds to, and suppress signals from, activated receptors tyrosine kinases, including the insulin (INSR) and insulin-like growth factor (IGF1R) receptors. The inhibitory effect can be achieved by 2 mechanisms: interference with the signaling pathway and increased receptor degradation. Delays and reduces AKT1 phosphorylation in response to insulin stimulation. Blocks association between INSR and IRS1 and IRS2 and prevents insulin-stimulated IRS1 and IRS2 tyrosine phosphorylation. Recruits NEDD4 to IGF1R, leading to IGF1R ubiquitination, increased internalization and degradation by both the proteasomal and lysosomal pathways. May play a role in mediating insulin-stimulated ubiquitination of INSR, leading to proteasomal degradation. Negatively regulates Wnt signaling by interacting with LRP6 intracellular portion and interfering with the binding of AXIN1 to LRP6. Positive regulator of the KDR/VEGFR-2 signaling pathway. May inhibit NEDD4-mediated degradation of KDR/VEGFR-2.

#### Cellular Location

Cytoplasm. Note=When complexed with NEDD4 and IGF1R, follows IGF1R internalization, remaining associated with early endosomes. Uncouples from IGF1R-containing endosomes before the sorting of the receptor to the lysosomal compartment (By similarity).

#### Tissue Location

Widely expressed in fetal and adult tissues, including fetal and postnatal liver, lung, kidney, skeletal muscle, heart, spleen, skin and brain.

### GRB10 Antibody (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### GRB10 Antibody (internal region) - Images



AF2737a (1 µg/ml) staining of HeLa lysate (35 µg protein in RIPA buffer). Primary incubation was

1 hour. Detected by chemiluminescence.

#### **GRB10 Antibody (internal region) - Background**

This antibody is expected to recognise the reported isoforms ( NP\_005302.3; NP\_001001549.1)

#### **GRB10 Antibody (internal region) - References**

Up-regulation of growth factor receptor-bound protein 10 in cervical squamous cell carcinoma.  
Okino K, Konishi H, Doi D, Yoneyama K, Ota Y, Jin E, Kawanami O, Takeshita T. Oncol Rep. 2005 Jun;13(6):1069-74. PMID: 15870923