

# CDC6 Antibody (internal region)

Peptide-affinity purified goat antibody Catalog # AF3170a

#### Specification

## CDC6 Antibody (internal region) - Product Information

Application Primary Accession Other Accession Reactivity Host Clonality Concentration Isotype Calculated MW

WB, E <u>Q99741</u> <u>NP\_001245.1</u>, <u>990</u> Human Goat Polyclonal 0.5 mg/ml IgG 62720

#### CDC6 Antibody (internal region) - Additional Information

Gene ID 990

**Other Names** Cell division control protein 6 homolog, CDC6-related protein, Cdc18-related protein, HsCdc18, p62(cdc6), HsCDC6, CDC6, CDC18L

Dilution WB~~1:1000 E~~N/A

**Format** 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

**Storage** Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions** CDC6 Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

## CDC6 Antibody (internal region) - Protein Information

Name CDC6

Synonyms CDC18L

**Function** Involved in the initiation of DNA replication. Also participates in checkpoint controls that ensure



DNA replication is completed before mitosis is initiated.

**Cellular Location** 

Nucleus. Cytoplasm Note=The protein is nuclear in G1 and cytoplasmic in S-phase cells (PubMed:9566895).

#### CDC6 Antibody (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

## CDC6 Antibody (internal region) - Images

_	250kDa 150kDa 100kDa 75kDa	
	50kDa	
	37kDa	
	25kDa	
	20kDa	
	15kDa	

AF3170a (0.01  $\mu$ g/ml) staining of nuclear Jurkat lysate (35  $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

#### CDC6 Antibody (internal region) - References

ATR (AT mutated Rad3 related) activity stabilizes Cdc6 and delays G2/M-phase entry during hydroxyurea-induced S-phase arrest of HeLa cells. Liu L, Choi JH, Yim H, Choi JS, Park BD, Cho SJ, Lee SK, The international journal of biochemistry & cell biology 2009 Jun 41 (6): 1410-20. PMID: 19154794