

ALDH3A1 Antibody (C-Term)

Peptide-affinity purified goat antibody Catalog # AF3239a

Specification

ALDH3A1 Antibody (C-Term) - Product Information

Application WB, IHC, E
Primary Accession P30838
Other Accession NP_000682.3, 218
Reactivity Human, Mouse

Reactivity
Host
Clonality
Concentration
Human, Mouse
Goat
Polyclonal
0.5 mg/ml

Isotype IgG Calculated MW 50395

ALDH3A1 Antibody (C-Term) - Additional Information

Gene ID 218

Other Names

Aldehyde dehydrogenase, dimeric NADP-preferring, 1.2.1.5, ALDHIII, Aldehyde dehydrogenase 3, Aldehyde dehydrogenase family 3 member A1, ALDH3A1, ALDH3

Dilution

WB~~1:1000 IHC~~1:100~500

E~~N/A

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ALDH3A1 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

ALDH3A1 Antibody (C-Term) - Protein Information

Name ALDH3A1

Synonyms ALDH3

Function



ALDHs play a major role in the detoxification of alcohol- derived acetaldehyde (Probable). They are involved in the metabolism of corticosteroids, biogenic amines, neurotransmitters, and lipid peroxidation (Probable). Oxidizes medium and long chain aldehydes into non-toxic fatty acids (PubMed:1737758). Preferentially oxidizes aromatic aldehyde substrates (PubMed:1737758). Comprises about 50 percent of corneal epithelial soluble proteins (By similarity). May play a role in preventing

Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:P47739}.

corneal damage caused by ultraviolet light (By similarity).

Tissue Location


High levels in stomach, esophagus and lung; low level in the liver and kidney

ALDH3A1 Antibody (C-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

ALDH3A1 Antibody (C-Term) - Images



AF3239a (0.01 μ g/ml) staining of Mouse Eye lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.





AF3239a (3.8 μ g/ml) staining of paraffin embedded Human Lung. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.

ALDH3A1 Antibody (C-Term) - Background

Reported variants represent identical protein: NP_001128639.1, NP_000682.3, NP_001128640.1

ALDH3A1 Antibody (C-Term) - References

The oxidation status of ALDH3A1 in human saliva and its correlation with antioxidant capacity measured by ORAC method. Bogucka M, Giebu?towicz J, Zawada K, Wroczy?ski P, Wierzchowski J, Pietrzak M, Piekarczyk P, Romanowska K. Acta poloniae pharmaceutica 66 (5): 477-82. PMID: 19894643