

**DBH (aa437-448) Antibody (internal region)**  
**Peptide-affinity purified goat antibody**  
**Catalog # AF3702a****Specification**

---

**DBH (aa437-448) Antibody (internal region) - Product Information**

|                   |                                                                                                                  |
|-------------------|------------------------------------------------------------------------------------------------------------------|
| Application       | WB, E                                                                                                            |
| Primary Accession | <a href="#">P09172</a>                                                                                           |
| Other Accession   | <a href="#">NP_000778.3</a> , <a href="#">1621</a> , <a href="#">13166 (mouse)</a> , <a href="#">25699 (rat)</a> |
| Reactivity        | Human                                                                                                            |
| Predicted         | Mouse, Rat, Pig                                                                                                  |
| Host              | Goat                                                                                                             |
| Clonality         | Polyclonal                                                                                                       |
| Concentration     | 0.5 mg/ml                                                                                                        |
| Isotype           | IgG                                                                                                              |
| Calculated MW     | 69065                                                                                                            |

**DBH (aa437-448) Antibody (internal region) - Additional Information****Gene ID** 1621**Other Names**

Dopamine beta-hydroxylase, 1.14.17.1, Dopamine beta-monooxygenase, Soluble dopamine beta-hydroxylase, DBH

**Dilution**

WB~~1:1000

E~~N/A

**Format**

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

DBH (aa437-448) Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

**DBH (aa437-448) Antibody (internal region) - Protein Information****Name** DBH**Function**

Catalyzes the hydroxylation of dopamine to noradrenaline (also known as norepinephrine), and is

thus vital for regulation of these neurotransmitters.

#### Cellular Location


[Soluble dopamine beta-hydroxylase]: Cytoplasmic vesicle, secretory vesicle lumen Cytoplasmic vesicle, secretory vesicle, chromaffin granule lumen. Secreted

#### DBH (aa437-448) Antibody (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### DBH (aa437-448) Antibody (internal region) - Images



AF3702a (2 µg/ml) staining of Jurkat lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

#### DBH (aa437-448) Antibody (internal region) - References

Norepinephrine deficiency is caused by combined abnormal mRNA processing and defective protein trafficking of dopamine beta-hydroxylase. Kim CH, Leung A, Huh YH, Yang E, Kim DJ, Leblanc P, Ryu H, Kim K, Kim DW, Garland EM, Raj SR, Biaggioni I, Robertson D, Kim KS. J Biol Chem. 2011 Mar 18;286(11):9196-204. Epub 2011 Jan 5. PMID: 21209083