

**fibrillarin / FBL (aa99-110) Antibody (internal region)**  
**Peptide-affinity purified goat antibody**  
**Catalog # AF3811a****Specification**

---

**fibrillarin / FBL (aa99-110) Antibody (internal region) - Product Information**

|                   |                                                                                                                   |
|-------------------|-------------------------------------------------------------------------------------------------------------------|
| Application       | WB, E                                                                                                             |
| Primary Accession | <a href="#">P22087</a>                                                                                            |
| Other Accession   | <a href="#">NP_001427.2</a> , <a href="#">2091</a> , <a href="#">14113 (mouse)</a> , <a href="#">292747 (rat)</a> |
| Reactivity        | Human                                                                                                             |
| Predicted         | Mouse, Rat, Pig, Dog                                                                                              |
| Host              | Goat                                                                                                              |
| Clonality         | Polyclonal                                                                                                        |
| Concentration     | 0.5 mg/ml                                                                                                         |
| Isotype           | IgG                                                                                                               |
| Calculated MW     | 33784                                                                                                             |

**fibrillarin / FBL (aa99-110) Antibody (internal region) - Additional Information****Gene ID** 2091**Other Names**

rRNA 2'-O-methyltransferase fibrillarin, 2.1.1.-, 34 kDa nucleolar scleroderma antigen, Histone-glutamine methyltransferase, FBL, FIB1, FLRN

**Dilution**

WB~~1:1000

E~~N/A

**Format**

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

fibrillarin / FBL (aa99-110) Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

**fibrillarin / FBL (aa99-110) Antibody (internal region) - Protein Information****Name** FBL ([HGNC:3599](#))**Synonyms** FIB1, FLRN

### Function

S-adenosyl-L-methionine-dependent methyltransferase that has the ability to methylate both RNAs and proteins (PubMed:<a href="http://www.uniprot.org/citations/24352239" target="\_blank">24352239</a>, PubMed:<a href="http://www.uniprot.org/citations/30540930" target="\_blank">30540930</a>, PubMed:<a href="http://www.uniprot.org/citations/32017898" target="\_blank">32017898</a>). Involved in pre-rRNA processing by catalyzing the site-specific 2'-hydroxyl methylation of ribose moieties in pre-ribosomal RNA (PubMed:<a href="http://www.uniprot.org/citations/30540930" target="\_blank">30540930</a>). Site specificity is provided by a guide RNA that base pairs with the substrate (By similarity). Methylation occurs at a characteristic distance from the sequence involved in base pairing with the guide RNA (By similarity). Probably catalyzes 2'-O-methylation of U6 snRNAs in box C/D RNP complexes (PubMed:<a href="http://www.uniprot.org/citations/32017898" target="\_blank">32017898</a>). U6 snRNA 2'-O-methylation is required for mRNA splicing fidelity (PubMed:<a href="http://www.uniprot.org/citations/32017898" target="\_blank">32017898</a>). Also acts as a protein methyltransferase by mediating methylation of 'Gln-105' of histone H2A (H2AQ104me), a modification that impairs binding of the FACT complex and is specifically present at 35S ribosomal DNA locus (PubMed:<a href="http://www.uniprot.org/citations/24352239" target="\_blank">24352239</a>, PubMed:<a href="http://www.uniprot.org/citations/30540930" target="\_blank">30540930</a>). Part of the small subunit (SSU) processome, first precursor of the small eukaryotic ribosomal subunit. During the assembly of the SSU processome in the nucleolus, many ribosome biogenesis factors, an RNA chaperone and ribosomal proteins associate with the nascent pre-rRNA and work in concert to generate RNA folding, modifications, rearrangements and cleavage as well as targeted degradation of pre-ribosomal RNA by the RNA exosome (PubMed:<a href="http://www.uniprot.org/citations/34516797" target="\_blank">34516797</a>).

### Cellular Location


Nucleus, nucleolus. Nucleus, nucleoplasm {ECO:0000250|UniProtKB:P35550}. Note=Fibrillar region of the nucleolus


### fibrillarin / FBL (aa99-110) Antibody (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### fibrillarin / FBL (aa99-110) Antibody (internal region) - Images





AF3811a (0.1 µg/ml) staining of HEK293 lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

#### **fibrillarin / FBL (aa99-110) Antibody (internal region) - References**

Association of guanine nucleotide-exchange protein BIG1 in HepG2 cell nuclei with nucleolin, U3 snoRNA, and fibrillarin. Padilla PI, Uhart M, Pacheco-Rodriguez G, Peculis BA, Moss J, Vaughan M. Proc Natl Acad Sci U S A. 2008 Mar 4;105(9):3357-61. Epub 2008 Feb 21. PMID: 18292223