

AMACR Antibody (C-Term)
Peptide-affinity purified goat antibody
Catalog # AF4124a**Specification**

AMACR Antibody (C-Term) - Product Information

Application	WB, Pep-ELISA
Primary Accession	O9UHK6
Other Accession	NP_055139.4 , NP_001161067.1 , 23600
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	42387

AMACR Antibody (C-Term) - Additional Information**Gene ID** 23600**Other Names**

Alpha-methylacyl-CoA racemase, 5.1.99.4, 2-methylacyl-CoA racemase, AMACR

Dilution

WB~~1:1000

Pep-ELISA~~N/A

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

AMACR Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

AMACR Antibody (C-Term) - Protein Information**Name** AMACR**Function**

Catalyzes the interconversion of (R)- and (S)-stereoisomers of alpha-methyl-branched-chain fatty acyl-CoA esters (PubMed:10655068, PubMed:11060359, PubMed:7649182)

target="_blank">7649182). Acts only on coenzyme A thioesters, not on free fatty acids, and accepts as substrates a wide range of alpha-methylacyl-CoAs, including pristanoyl-CoA, trihydroxycoprostanoyl-CoA (an intermediate in bile acid synthesis), and arylpropionic acids like the anti-inflammatory drug ibuprofen (2- (4-isobutylphenyl)propionic acid) but neither 3-methyl-branched nor linear-chain acyl-CoAs (PubMed:10655068, PubMed:11060359, PubMed:7649182).

Cellular Location


Peroxisome. Mitochondrion

AMACR Antibody (C-Term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


AMACR Antibody (C-Term) - Images



AF4124a (0.5 µg/ml) staining of Human Kidney lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



EB12538 (0.5µg/ml) staining of CaCo-2 (A) and LNCaP (B) cell lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.



EB12538 (0.5µg/ml) staining of Human Kidney (A) and Liver (B) lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.

AMACR Antibody (C-Term) - Background

This antibody is expected to recognize isoform 1 (NP_055139.4) and isoform 3 (NP_001161067.1).

AMACR Antibody (C-Term) - References

High-throughput screen identifies novel inhibitors of cancer biomarker a-methylacyl coenzyme A racemase (AMACR/P504S). Wilson BA, Wang H, Nacev BA, Mease RC, Liu JO, Pomper MG, Isaacs WB. Molecular cancer therapeutics 2011 May 10 (5): 825-38. PMID: 21441411