

**Goat Anti-TICAM-1 / TRIF Antibody (internal region)**  
**Purified Goat Polyclonal Antibody**  
**Catalog # AF4229a****Specification**

---

**Goat Anti-TICAM-1 / TRIF Antibody (internal region) - Product Information**

|                   |                                                                                           |
|-------------------|-------------------------------------------------------------------------------------------|
| Application       | WB, E                                                                                     |
| Primary Accession | <a href="#">Q8IUC6</a>                                                                    |
| Other Accession   | <a href="#">106759(mouse)</a> , <a href="#">363328(rat)</a> , <a href="#">NP_891549.1</a> |
| Reactivity        | Human                                                                                     |
| Predicted         | Human, Mouse, Rat, Pig, Dog                                                               |
| Host              | Goat                                                                                      |
| Clonality         | Polyclonal                                                                                |
| Concentration     | 0.5                                                                                       |
| Calculated MW     | 76422                                                                                     |

**Goat Anti-TICAM-1 / TRIF Antibody (internal region) - Additional Information****Gene ID** 148022**Other Names**

TICAM1; toll-like receptor adaptor molecule 1; PRVTIRB; TICAM-1; TRIF; TIR domain containing adaptor inducing interferon-beta; TIR domain-containing adapter molecule 1; TIR domain-containing adapter protein inducing IFN-beta; proline-rich, vinculin and TIR domain-containing protein B; putative NF-kappa-B-activating protein 502H; toll-interleukin-1 receptor domain-containing adapter protein inducing interferon beta

**Dilution**

WB~~1:1000  
E~~N/A

**Format**

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.

**Immunogen**

Peptide with sequence PDGATFCEDFQVP, from the internal region of the protein sequence according to NP\_891549.1.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Goat Anti-TICAM-1 / TRIF Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

**Goat Anti-TICAM-1 / TRIF Antibody (internal region) - Protein Information**

**Name** TICAM1**Synonyms** PRVTIRB, TRIF**Function**

Involved in innate immunity against invading pathogens. Adapter used by TLR3, TLR4 (through TICAM2) and TLR5 to mediate NF- $\kappa$ B and interferon-regulatory factor (IRF) activation, and to induce apoptosis (PubMed:<a href="http://www.uniprot.org/citations/12471095" target="\_blank">12471095</a>, PubMed:<a href="http://www.uniprot.org/citations/12539043" target="\_blank">12539043</a>, PubMed:<a href="http://www.uniprot.org/citations/14739303" target="\_blank">14739303</a>, PubMed:<a href="http://www.uniprot.org/citations/28747347" target="\_blank">28747347</a>, PubMed:<a href="http://www.uniprot.org/citations/35215908" target="\_blank">35215908</a>). Ligand binding to these receptors results in TRIF recruitment through its TIR domain (PubMed:<a href="http://www.uniprot.org/citations/12471095" target="\_blank">12471095</a>, PubMed:<a href="http://www.uniprot.org/citations/12539043" target="\_blank">12539043</a>, PubMed:<a href="http://www.uniprot.org/citations/14739303" target="\_blank">14739303</a>). Distinct protein-interaction motifs allow recruitment of the effector proteins TBK1, TRAF6 and RIPK1, which in turn, lead to the activation of transcription factors IRF3 and IRF7, NF- $\kappa$ B and FADD respectively (PubMed:<a href="http://www.uniprot.org/citations/12471095" target="\_blank">12471095</a>, PubMed:<a href="http://www.uniprot.org/citations/12539043" target="\_blank">12539043</a>, PubMed:<a href="http://www.uniprot.org/citations/14739303" target="\_blank">14739303</a>). Phosphorylation by TBK1 on the pLxIS motif leads to recruitment and subsequent activation of the transcription factor IRF3 to induce expression of type I interferon and exert a potent immunity against invading pathogens (PubMed:<a href="http://www.uniprot.org/citations/25636800" target="\_blank">25636800</a>). Component of a multi- helicase-TICAM1 complex that acts as a cytoplasmic sensor of viral double-stranded RNA (dsRNA) and plays a role in the activation of a cascade of antiviral responses including the induction of pro- inflammatory cytokines (By similarity).

**Cellular Location**

Cytoplasmic vesicle, autophagosome. Cytoplasm, cytosol {ECO:0000250|UniProtKB:Q80UF7}. Mitochondrion {ECO:0000250|UniProtKB:Q80UF7}. Note=Colocalizes with UBQLN1 in the autophagosome (PubMed:21695056). Colocalizes in the cytosol with DDX1, DDX21 and DHX36. Colocalizes in the mitochondria with DDX1 and poly(I:C) RNA ligand. The multi-helicase-TICAM1 complex may translocate to the mitochondria upon poly(I:C) RNA ligand stimulation (By similarity). {ECO:0000250|UniProtKB:Q80UF7, ECO:0000269|PubMed:21695056}

**Tissue Location**


Ubiquitously expressed but with higher levels in liver.

**Goat Anti-TICAM-1 / TRIF Antibody (internal region) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Goat Anti-TICAM-1 / TRIF Antibody (internal region) - Images**



AF4229a (2 µg/ml) staining of Daudi lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

**Goat Anti-TICAM-1 / TRIF Antibody (internal region) - References**

Selective TRIF-dependent signaling by a synthetic toll-like receptor 4 agonist. Bowen WS, Minns LA, Johnson DA, Mitchell TC, Hutton MM, Evans JT. Science signaling 2012 Feb 5 (211): ra13.