

**Goat Anti-NONO / p54NRB Antibody (C Terminus), biotinylated**  
**Catalog # AF4288a****Specification****Goat Anti-NONO / p54NRB Antibody (C Terminus), biotinylated - Product Information**

Application	WB, E
Primary Accession	<a href="#">Q15233</a>
Other Accession	<a href="#">NP_031389.3</a> , <a href="#">317259</a> , <a href="#">53610</a> , <a href="#">4841</a>
Reactivity	Human, Mouse
Predicted	Human, Mouse, Rat, Dog
Host	Goat
Isotype	IgG
Calculated MW	54232

**Goat Anti-NONO / p54NRB Antibody (C Terminus), biotinylated - Additional Information****Gene ID** 4841**Other Names**

Non-POU domain-containing octamer-binding protein, NonO protein, 54 kDa nuclear RNA- and DNA-binding protein, 55 kDa nuclear protein, DNA-binding p52/p100 complex, 52 kDa subunit, NMT55, p54(nrb), p54nrb, NONO, NRB54

**Dilution**

WB~~1:1000

E~~N/A

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Goat Anti-NONO / p54NRB Antibody (C Terminus), biotinylated is for research use only and not for use in diagnostic or therapeutic procedures.

**Goat Anti-NONO / p54NRB Antibody (C Terminus), biotinylated - Protein Information****Name** NONO {ECO:0000303|PubMed:9393982, ECO:0000312|HGNC:HGNC:7871}**Function**

DNA- and RNA binding protein, involved in several nuclear processes (PubMed:<a href="http://www.uniprot.org/citations/11525732" target="\_blank">11525732</a>, PubMed:<a href="http://www.uniprot.org/citations/12403470" target="\_blank">12403470</a>, PubMed:<a href="http://www.uniprot.org/citations/26571461" target="\_blank">26571461</a>). Binds the conventional octamer sequence in double-stranded DNA (PubMed:<a href="http://www.uniprot.org/citations/11525732" target="\_blank">11525732</a>, PubMed:<a href="http://www.uniprot.org/citations/12403470" target="\_blank">12403470</a>, PubMed:<a href="http://www.uniprot.org/citations/26571461" target="\_blank">26571461</a>).

<http://www.uniprot.org/citations/26571461> target="\_blank">26571461</a>). Also binds single-stranded DNA and RNA at a site independent of the duplex site (PubMed:<a href="http://www.uniprot.org/citations/11525732" target="\_blank">11525732</a>, PubMed:<a href="http://www.uniprot.org/citations/12403470" target="\_blank">12403470</a>, PubMed:<a href="http://www.uniprot.org/citations/26571461" target="\_blank">26571461</a>). Involved in pre-mRNA splicing, probably as a heterodimer with SFPQ (PubMed:<a href="http://www.uniprot.org/citations/11525732" target="\_blank">11525732</a>, PubMed:<a href="http://www.uniprot.org/citations/12403470" target="\_blank">12403470</a>, PubMed:<a href="http://www.uniprot.org/citations/26571461" target="\_blank">26571461</a>). Interacts with U5 snRNA, probably by binding to a purine-rich sequence located on the 3' side of U5 snRNA stem 1b (PubMed:<a href="http://www.uniprot.org/citations/12403470" target="\_blank">12403470</a>). Together with PSPC1, required for the formation of nuclear paraspeckles (PubMed:<a href="http://www.uniprot.org/citations/22416126" target="\_blank">22416126</a>). The SFPQ-NONO heteromer associated with MATR3 may play a role in nuclear retention of defective RNAs (PubMed:<a href="http://www.uniprot.org/citations/11525732" target="\_blank">11525732</a>). The SFPQ-NONO heteromer may be involved in DNA unwinding by modulating the function of topoisomerase I/TOPI (PubMed:<a href="http://www.uniprot.org/citations/10858305" target="\_blank">10858305</a>). The SFPQ-NONO heteromer may be involved in DNA non-homologous end joining (NHEJ) required for double-strand break repair and V(D)J recombination and may stabilize paired DNA ends (PubMed:<a href="http://www.uniprot.org/citations/15590677" target="\_blank">15590677</a>). In vitro, the complex strongly stimulates DNA end joining, binds directly to the DNA substrates and cooperates with the Ku70/G22P1-Ku80/XRCC5 (Ku) dimer to establish a functional preligation complex (PubMed:<a href="http://www.uniprot.org/citations/15590677" target="\_blank">15590677</a>). NONO is involved in transcriptional regulation. The SFPQ-NONO-NR5A1 complex binds to the CYP17 promoter and regulates basal and cAMP-dependent transcriptional activity (PubMed:<a href="http://www.uniprot.org/citations/11897684" target="\_blank">11897684</a>). NONO binds to an enhancer element in long terminal repeats of endogenous intracisternal A particles (IAPs) and activates transcription (By similarity). Regulates the circadian clock by repressing the transcriptional activator activity of the CLOCK-BMAL1 heterodimer (By similarity). Important for the functional organization of GABAergic synapses (By similarity). Plays a specific and important role in the regulation of synaptic RNAs and GPHN/gephyrin scaffold structure, through the regulation of GABRA2 transcript (By similarity). Plays a key role during neuronal differentiation by recruiting TET1 to genomic loci and thereby regulating 5-hydroxymethylcytosine levels (By similarity). Plays a role in the regulation of DNA virus-mediated innate immune response by assembling into the HDP-RNP complex, a complex that serves as a platform for IRF3 phosphorylation and subsequent innate immune response activation through the cGAS-STING pathway (PubMed:<a href="http://www.uniprot.org/citations/28712728" target="\_blank">28712728</a>, PubMed:<a href="http://www.uniprot.org/citations/30270045" target="\_blank">30270045</a>). Promotes activation of the cGAS-STING pathway in response to HIV-2 infection: acts by interacting with HIV-2 Capsid protein p24, thereby promoting detection of viral DNA by CGAS, leading to CGAS-mediated immune activation (PubMed:<a href="http://www.uniprot.org/citations/30270045" target="\_blank">30270045</a>). In contrast, the weak interaction with HIV-1 Capsid protein p24 does not allow activation of the cGAS-STING pathway (PubMed:<a href="http://www.uniprot.org/citations/30270045" target="\_blank">30270045</a>).

### Cellular Location

Nucleus. Nucleus, nucleolus. Nucleus speckle. Chromosome {ECO:0000250|UniProtKB:Q99K48}. Note=Detected in punctate subnuclear structures often located adjacent to splicing speckles, called paraspeckles.

### Tissue Location

Heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Also found in a number of breast tumor cell lines.

**Goat Anti-NONO / p54NRB Antibody (C Terminus), biotinylated - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Goat Anti-NONO / p54NRB Antibody (C Terminus), biotinylated - Images**

Biotinylated AF4288a (0.1 µg/ml) staining of mouse brain lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.