

Goat anti-ACADM, Biotinylated Antibody
Peptide-affinity purified goat antibody
Catalog # AF4364a**Specification**

Goat anti-ACADM, Biotinylated Antibody - Product Information

Application	WB, IHC, Pep-ELISA
Primary Accession	P11310
Other Accession	NP_000007.1 , NP_001120800.1 , NP_001272971.1 , NP_001272972.1 , NP_001272973.1
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Calculated MW	46588

Goat anti-ACADM, Biotinylated Antibody - Additional Information**Gene ID** 34**Other Names**ACADM; acyl-CoA dehydrogenase, C-4 to C-12 straight chain; ACAD1; MCAD; MCADH;
acyl-Coenzyme A dehydrogenase, C-4 to C-12 straight chain**Dilution**WB~~1:1000
IHC~~1:100~500
Pep-ELISA~~N/A**Format**Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.
Aliquot and store at -20°C. Minimize freezing and thawing.**Immunogen**

This antibody is expected to recognise both reported isoforms (NP_000007.1; NP_001120800.1).

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat anti-ACADM, Biotinylated Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat anti-ACADM, Biotinylated Antibody - Protein Information**Name** ACADM ([HGNC:89](#))

Function

Medium-chain specific acyl-CoA dehydrogenase is one of the acyl-CoA dehydrogenases that catalyze the first step of mitochondrial fatty acid beta-oxidation, an aerobic process breaking down fatty acids into acetyl-CoA and allowing the production of energy from fats (PubMed:1970566, PubMed:21237683, PubMed:2251268, PubMed:8823175). The first step of fatty acid beta-oxidation consists in the removal of one hydrogen from C-2 and C-3 of the straight-chain fatty acyl-CoA thioester, resulting in the formation of trans-2-enoyl-CoA (PubMed:2251268). Electron transfer flavoprotein (ETF) is the electron acceptor that transfers electrons to the main mitochondrial respiratory chain via ETF-ubiquinone oxidoreductase (ETF dehydrogenase) (PubMed:15159392, PubMed:25416781). Among the different mitochondrial acyl-CoA dehydrogenases, medium-chain specific acyl-CoA dehydrogenase acts specifically on acyl-CoAs with saturated 6 to 12 carbons long primary chains (PubMed:1970566, PubMed:21237683, PubMed:2251268, PubMed:8823175).

Cellular Location

Mitochondrion matrix

Goat anti-ACADM, Biotinylated Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat anti-ACADM, Biotinylated Antibody - Images