

**Goat anti-RSK2 Antibody**  
**Peptide-affinity purified goat antibody**  
**Catalog # AF4543a****Specification**

---

**Goat anti-RSK2 Antibody - Product Information**

|                   |                             |
|-------------------|-----------------------------|
| Application       | IHC, Pep-ELISA              |
| Primary Accession | <a href="#">P51812</a>      |
| Other Accession   | <a href="#">NP_004577.1</a> |
| Reactivity        | Human, Mouse, Rat           |
| Host              | Goat                        |
| Clonality         | Polyclonal                  |
| Calculated MW     | 83736                       |

**Goat anti-RSK2 Antibody - Additional Information****Gene ID** 6197**Other Names**

RPS6KA3; ribosomal protein S6 kinase, 90kDa, polypeptide 3 ; HGNC:10432; HU-3; ISPK-1; MAPKAPK1B; MRX19; RSK2; S6K-alpha3; p90-RSK2; pp90RSK2; RP11-393H10.3; CLS; RSK; insulin-stimulated protein kinase 1; mental retardation, X-linked 19; ribosomal protein

**Dilution**

IHC~~1:100~500

Pep-ELISA~~N/A

**Format**

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Goat anti-RSK2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Goat anti-RSK2 Antibody - Protein Information****Name** RPS6KA3**Synonyms** ISPK1, MAPKAPK1B, RSK2**Function**

Serine/threonine-protein kinase that acts downstream of ERK (MAPK1/ERK2 and MAPK3/ERK1)

signaling and mediates mitogenic and stress-induced activation of the transcription factors CREB1, ETV1/ER81 and NR4A1/NUR77, regulates translation through RPS6 and EIF4B phosphorylation, and mediates cellular proliferation, survival, and differentiation by modulating mTOR signaling and repressing pro- apoptotic function of BAD and DAPK1 (PubMed:<a href="http://www.uniprot.org/citations/16213824" target="\_blank">16213824</a>, PubMed:<a href="http://www.uniprot.org/citations/16223362" target="\_blank">16223362</a>, PubMed:<a href="http://www.uniprot.org/citations/17360704" target="\_blank">17360704</a>, PubMed:<a href="http://www.uniprot.org/citations/9770464" target="\_blank">9770464</a>). In fibroblast, is required for EGF- stimulated phosphorylation of CREB1 and histone H3 at 'Ser-10', which results in the subsequent transcriptional activation of several immediate-early genes (PubMed:<a href="http://www.uniprot.org/citations/10436156" target="\_blank">10436156</a>, PubMed:<a href="http://www.uniprot.org/citations/9770464" target="\_blank">9770464</a>). In response to mitogenic stimulation (EGF and PMA), phosphorylates and activates NR4A1/NUR77 and ETV1/ER81 transcription factors and the cofactor CREBBP (PubMed:<a href="http://www.uniprot.org/citations/16223362" target="\_blank">16223362</a>). Upon insulin-derived signal, acts indirectly on the transcription regulation of several genes by phosphorylating GSK3B at 'Ser-9' and inhibiting its activity (PubMed:<a href="http://www.uniprot.org/citations/8250835" target="\_blank">8250835</a>). Phosphorylates RPS6 in response to serum or EGF via an mTOR-independent mechanism and promotes translation initiation by facilitating assembly of the preinitiation complex (PubMed:<a href="http://www.uniprot.org/citations/17360704" target="\_blank">17360704</a>). In response to insulin, phosphorylates EIF4B, enhancing EIF4B affinity for the EIF3 complex and stimulating cap-dependent translation (PubMed:<a href="http://www.uniprot.org/citations/18508509" target="\_blank">18508509</a>, PubMed:<a href="http://www.uniprot.org/citations/18813292" target="\_blank">18813292</a>). Is involved in the mTOR nutrient-sensing pathway by directly phosphorylating TSC2 at 'Ser-1798', which potently inhibits TSC2 ability to suppress mTOR signaling, and mediates phosphorylation of RPTOR, which regulates mTORC1 activity and may promote rapamycin- sensitive signaling independently of the PI3K/AKT pathway (PubMed:<a href="http://www.uniprot.org/citations/18722121" target="\_blank">18722121</a>). Mediates cell survival by phosphorylating the pro- apoptotic proteins BAD and DAPK1 and suppressing their pro-apoptotic function (PubMed:<a href="http://www.uniprot.org/citations/16213824" target="\_blank">16213824</a>). Promotes the survival of hepatic stellate cells by phosphorylating CEBPB in response to the hepatotoxin carbon tetrachloride (CCl4) (PubMed:<a href="http://www.uniprot.org/citations/18508509" target="\_blank">18508509</a>, PubMed:<a href="http://www.uniprot.org/citations/18813292" target="\_blank">18813292</a>). Is involved in cell cycle regulation by phosphorylating the CDK inhibitor CDKN1B, which promotes CDKN1B association with 14-3-3 proteins and prevents its translocation to the nucleus and inhibition of G1 progression (By similarity). In LPS-stimulated dendritic cells, is involved in TLR4- induced macropinocytosis, and in myeloma cells, acts as effector of FGFR3-mediated transformation signaling, after direct phosphorylation at Tyr-529 by FGFR3 (By similarity). Negatively regulates EGF-induced MAPK1/3 phosphorylation via phosphorylation of SOS1 (By similarity). Phosphorylates SOS1 at 'Ser-1134' and 'Ser-1161' that create YWHAB and YWHAЕ binding sites and which contribute to the negative regulation of MAPK1/3 phosphorylation (By similarity). Phosphorylates EPHA2 at 'Ser- 897', the RPS6KA-EPHA2 signaling pathway controls cell migration (PubMed:<a href="http://www.uniprot.org/citations/26158630" target="\_blank">26158630</a>). Acts as a regulator of osteoblast differentiation by mediating phosphorylation of ATF4, thereby promoting ATF4 transactivation activity (By similarity).

**Cellular Location**

Nucleus. Cytoplasm

**Tissue Location**

Expressed in many tissues, highest levels in skeletal muscle

**Goat anti-RSK2 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Goat anti-RSK2 Antibody - Images**