

**KD-Validated Anti-Annexin A1 Rabbit Monoclonal Antibody**  
**Rabbit monoclonal antibody**  
**Catalog # AGI1975****Specification****KD-Validated Anti-Annexin A1 Rabbit Monoclonal Antibody - Product Information**

|                   |                                                                                                                                                                                                                              |
|-------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Application       | WB, FC, ICC                                                                                                                                                                                                                  |
| Primary Accession | <a href="#">P04083</a>                                                                                                                                                                                                       |
| Reactivity        | Rat, Human, Mouse                                                                                                                                                                                                            |
| Clonality         | Monoclonal                                                                                                                                                                                                                   |
| Isotype           | Rabbit IgG                                                                                                                                                                                                                   |
| Calculated MW     | Predicted, 39 kDa, observed, 33 kDa kDa                                                                                                                                                                                      |
| Gene Name         | ANXA1                                                                                                                                                                                                                        |
| Aliases           | ANXA1; Annexin A1; ANX1; LPC1; Phospholipase A2 Inhibitory Protein; Chromobindin-9; Calpactin II; Calpactin-2; Annexin-1; Epididymis Secretory Sperm Binding Protein; Annexin I (Lipocortin I); Lipocortin I; Annexin I; P35 |
| Immunogen         | Recombinant protein of human Annexin A1                                                                                                                                                                                      |

**KD-Validated Anti-Annexin A1 Rabbit Monoclonal Antibody - Additional Information**

|                    |                                                                                                                                                                        |
|--------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Gene ID            | 301                                                                                                                                                                    |
| <b>Other Names</b> | Annexin A1, Annexin I, Annexin-1, Calpactin II, Calpactin-2, Chromobindin-9, Lipocortin I, Phospholipase A2 inhibitory protein, p35, Annexin Ac2-26, ANXA1, ANX1, LPC1 |

**KD-Validated Anti-Annexin A1 Rabbit Monoclonal Antibody - Protein Information****Name** ANXA1**Synonyms** ANX1, LPC1**Function**

Plays important roles in the innate immune response as effector of glucocorticoid-mediated responses and regulator of the inflammatory process. Has anti-inflammatory activity (PubMed:<a href="http://www.uniprot.org/citations/8425544" target="\_blank">8425544</a>). Plays a role in glucocorticoid-mediated down-regulation of the early phase of the inflammatory response (By similarity). Contributes to the adaptive immune response by enhancing signaling cascades that are triggered by T-cell activation, regulates differentiation and proliferation of activated T-cells (PubMed:<a href="http://www.uniprot.org/citations/17008549" target="\_blank">17008549</a>). Promotes the differentiation of T-cells into Th1 cells and negatively regulates differentiation into Th2 cells (PubMed:<a href="http://www.uniprot.org/citations/17008549" target="\_blank">17008549</a>). Has no effect on unstimulated T cells (PubMed:<a href="http://www.uniprot.org/citations/17008549" target="\_blank">17008549</a>). Negatively regulates hormone exocytosis via activation of the formyl peptide receptors and reorganization of the actin cytoskeleton (PubMed:<a href="http://www.uniprot.org/citations/19625660" target="\_blank">19625660</a>).

target="\_blank">19625660</a>). Has high affinity for Ca(2+) and can bind up to eight Ca(2+) ions (By similarity). Displays Ca(2+)-dependent binding to phospholipid membranes (PubMed:<a href="http://www.uniprot.org/citations/2532504" target="\_blank">2532504</a>, PubMed:<a href="http://www.uniprot.org/citations/8557678" target="\_blank">8557678</a>). Plays a role in the formation of phagocytic cups and phagosomes. Plays a role in phagocytosis by mediating the Ca(2+)-dependent interaction between phagosomes and the actin cytoskeleton (By similarity).

#### Cellular Location

Nucleus. Cytoplasm. Cell projection, cilium {ECO:0000250|UniProtKB:P46193}. Cell membrane. Membrane; Peripheral membrane protein. Endosome membrane {ECO:0000250|UniProtKB:P07150}; Peripheral membrane protein {ECO:0000250|UniProtKB:P07150}. Basolateral cell membrane {ECO:0000250|UniProtKB:P51662}. Apical cell membrane {ECO:0000250|UniProtKB:P10107}. Lateral cell membrane {ECO:0000250|UniProtKB:P10107}. Secreted. Secreted, extracellular space. Cell membrane; Peripheral membrane protein; Extracellular side. Secreted, extracellular exosome. Cytoplasmic vesicle, secretory vesicle lumen. Cell projection, phagocytic cup {ECO:0000250|UniProtKB:P10107}. Early endosome {ECO:0000250|UniProtKB:P19619}. Cytoplasmic vesicle membrane {ECO:0000250|UniProtKB:P19619}; Peripheral membrane protein {ECO:0000250|UniProtKB:P19619}. Note=Secreted, at least in part via exosomes and other secretory vesicles. Detected in exosomes and other extracellular vesicles (PubMed:25664854). Alternatively, the secretion is dependent on protein unfolding and facilitated by the cargo receptor TMED10; it results in the protein translocation from the cytoplasm into ERGIC (endoplasmic reticulum-Golgi intermediate compartment) followed by vesicle entry and secretion (PubMed:32272059). Detected in gelatinase granules in resting neutrophils (PubMed:10772777). Secretion is increased in response to wounding and inflammation (PubMed:25664854). Secretion is increased upon T-cell activation (PubMed:17008549). Neutrophil adhesion to endothelial cells stimulates secretion via gelatinase granules, but foreign particle phagocytosis has no effect (PubMed:10772777). Colocalizes with actin fibers at phagocytic cups (By similarity). Displays calcium-dependent binding to phospholipid membranes (PubMed:2532504, PubMed:8557678) {ECO:0000250|UniProtKB:P10107, ECO:0000269|PubMed:10772777, ECO:0000269|PubMed:17008549, ECO:0000269|PubMed:2532504, ECO:0000269|PubMed:25664854, ECO:0000269|PubMed:32272059, ECO:0000269|PubMed:8557678}

#### Tissue Location


Detected in resting neutrophils (PubMed:10772777). Detected in peripheral blood T-cells (PubMed:17008549). Detected in extracellular vesicles in blood serum from patients with inflammatory bowel disease, but not in serum from healthy donors (PubMed:25664854) Detected in placenta (at protein level) (PubMed:2532504). Detected in liver.


### KD-Validated Anti-Annexin A1 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### KD-Validated Anti-Annexin A1 Rabbit Monoclonal Antibody - Images







Western blotting analysis using anti-annexin A1 antibody (Cat#AGI1975). Total cell lysates (10 µg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-annexin A1 antibody (Cat#AGI1975, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.



Western blotting analysis using anti-annexin A1 antibody (Cat#AGI1975). Annexin A1 expression in wild-type (WT) and annexin A1 (ANXA1) shRNA knockdown (KD) HeLa cells with 20 µg of total cell lysates. Hsp90 α serves as a loading control. The blot was incubated with anti-annexin A1 antibody (Cat#AGI1975, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.



Flow cytometric analysis of Annexin A1 expression in HepG2 cells using anti-Annexin A1 antibody (Cat#AGI1975, 1:2,000). Green, isotype control; red, Annexin A1.



Immunocytochemical staining of HepG2 cells with anti-Annexin A1 antibody (Cat#AGI1975, 1:1,000). Nuclei were stained blue with DAPI; Annexin A1 was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser Intensity and smart gain: High. Scale bar, 20  $\mu$ m.