

KD-Validated Anti-Erb-b2 receptor tyrosine kinase 2 Rabbit Monoclonal Antibody
Rabbit monoclonal antibody
Catalog # AGI2053**Specification****KD-Validated Anti-Erb-b2 receptor tyrosine kinase 2 Rabbit Monoclonal Antibody - Product Information**

Application	WB, ICC
Primary Accession	P04626
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Isotype	Rabbit IgG
Calculated MW	Predicted, 138 kDa , observed, 185 kDa
Gene Name	KDa
Aliases	ERBB2 Erb-B2 Receptor Tyrosine Kinase 2; HER2; NEU; C-ERB-2; C-ERB2; MLN-19; HER-2; CD340; NGL; V-Erb-B2 Avian Erythroblastic Leukemia Viral Oncogene Homolog 2; Tyrosine Kinase-Type Cell Surface Receptor HER2; Neuro/Glioblastoma Derived Oncogene Homolog; Human Epidermal Growth Factor Receptor 2; Receptor Tyrosine-Protein Kinase ErbB-2; Metastatic Lymph Node Gene 19 Protein; Proto-Oncogene C-ErbB-2; Proto-Oncogene Neu; P185(ErbB2); EC 2.7.10.1; MLN 19; V-Erb-B2 Avian Erythroblastic Leukemia Viral Oncogene Homolog 2 (Neuro/Glioblastoma Derived Oncogene Homolog); V-Erb-B2 Erythroblastic Leukemia Viral Oncogene Homolog 2, Neuro/Glioblastoma Derived Oncogene Homolog 3V-Erb-B2 Avian Erythroblastic Leukemia Viral Oncoprotein; Neuroblastoma/Glioblastoma Derived Oncogene Homolog; Metastatic Lymph Node Gene 19; C-Erb B2/Neu Protein; CD340 Antigen; P185(ERBB2); HER-2/Neu; Herstatin; P185erbB2; EC 2.7.10; VSCN2; MLN19; TKR1
Immunogen	A synthesized peptide derived from human ErbB2 (HER2)

KD-Validated Anti-Erb-b2 receptor tyrosine kinase 2 Rabbit Monoclonal Antibody - Additional Information

Gene ID	2064
Other Names	
Receptor tyrosine-protein kinase erbB-2, 2.7.10.1, Metastatic lymph node gene 19 protein, MLN	

19, Proto-oncogene Neu, Proto-oncogene c-ErbB-2, Tyrosine kinase-type cell surface receptor
HER2, p185erbB2, CD340, ERBB2, HER2, MLN19, NEU, NGL

KD-Validated Anti-Erb-b2 receptor tyrosine kinase 2 Rabbit Monoclonal Antibody - Protein Information

Name ERBB2

Synonyms HER2, MLN19, NEU, NGL

Function

Protein tyrosine kinase that is part of several cell surface receptor complexes, but that apparently needs a coreceptor for ligand binding. Essential component of a neuregulin-receptor complex, although neuregulins do not interact with it alone. GP30 is a potential ligand for this receptor. Regulates outgrowth and stabilization of peripheral microtubules (MTs). Upon ERBB2 activation, the MEMO1-RHOA-DIAPH1 signaling pathway elicits the phosphorylation and thus the inhibition of GSK3B at cell membrane. This prevents the phosphorylation of APC and CLASP2, allowing its association with the cell membrane. In turn, membrane-bound APC allows the localization of MACF1 to the cell membrane, which is required for microtubule capture and stabilization.

Cellular Location

Cell membrane; Single-pass type I membrane protein. Cell projection, ruffle membrane; Single-pass type I membrane protein. Note=Internalized from the cell membrane in response to EGF stimulation. [Isoform 2]: Cytoplasm. Nucleus.

Tissue Location


Expressed in a variety of tumor tissues including primary breast tumors and tumors from small bowel, esophagus, kidney and mouth.


KD-Validated Anti-Erb-b2 receptor tyrosine kinase 2 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

KD-Validated Anti-Erb-b2 receptor tyrosine kinase 2 Rabbit Monoclonal Antibody - Images






Western blotting analysis using anti-Erb-b2 receptor tyrosine kinase 2 antibody (Cat#AGI2053). Total cell lysates (30 µg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-Erb-b2 receptor tyrosine kinase 2 antibody (Cat#AGI2053, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.



Western blotting analysis using anti-Erb-b2 receptor tyrosine kinase 2 antibody (Cat#AGI2053). Erb-b2 receptor tyrosine kinase 2 expression in wild type (WT) and erb-b2 receptor tyrosine kinase 2 shRNA knockdown (KD) 293T cells with 30 µg of total cell lysates. Hsp90 α serves as a loading control. The blot was incubated with anti-Erb-b2 receptor tyrosine kinase 2 antibody (Cat#AGI2053, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.



Immunocytochemical staining of HepG2 cells with Erb-b2 receptor tyrosine kinase 2 antibody (Cat#AGI2053, 1:1,000). Nuclei were stained blue with DAPI; Erb-b2 receptor tyrosine kinase 2 was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser Intensity and smart gain: Very low. Scale bar: 20 µm.