

KD-Validated Anti-Cullin 5 Rabbit Monoclonal Antibody
Rabbit monoclonal antibody
Catalog # AGI2249**Specification****KD-Validated Anti-Cullin 5 Rabbit Monoclonal Antibody - Product Information**

Application	WB
Primary Accession	Q93034
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Isotype	Rabbit IgG
Calculated MW	Predicted, 91 kDa; observed, 75 kDa kDa
Gene Name	CUL5
Aliases	CUL5; Cullin 5; VACM-1; Vasopressin-Activated Calcium-Mobilizing Receptor 1; Cullin-5; CUL-5; VACM1; Cullin-5 (Vasopressin-Activated Calcium-Mobilizing Receptor-1); Vasopressin-Activated Calcium-Mobilizing Receptor-1
Immunogen	Recombinant protein of human Cullin 5

KD-Validated Anti-Cullin 5 Rabbit Monoclonal Antibody - Additional Information

Gene ID	8065
Other Names	
Cullin-5, CUL-5, Vasopressin-activated calcium-mobilizing receptor 1, VACM-1, CUL5 {ECO:0000303 PubMed:10230407, ECO:0000312 HGNC:HGNC:2556}	

KD-Validated Anti-Cullin 5 Rabbit Monoclonal Antibody - Protein Information

Name CUL5 {ECO:0000303|PubMed:10230407, ECO:0000312|HGNC:HGNC:2556}

Function

Core component of multiple cullin-5-RING E3 ubiquitin-protein ligase complexes (ECS complexes, also named CRL5 complexes), which mediate the ubiquitination and subsequent proteasomal degradation of target proteins (PubMed:11384984, PubMed:15601820, PubMed:21199876, PubMed:21980433, PubMed:23897481, PubMed:25505247, PubMed:27910872, PubMed:32200094, PubMed:33268465, PubMed:35512830, PubMed:38418882). Acts as a scaffold protein that contributes to catalysis through

positioning of the substrate and the ubiquitin-conjugating enzyme (PubMed:11384984, PubMed:15601820, PubMed:33268465). The functional specificity of the E3 ubiquitin-protein ligase complex depends on the variable SOCS box-containing substrate recognition component (PubMed:11384984, PubMed:15601820, PubMed:33268465). Acts as a key regulator of neuron positioning during cortex development: component of various SOCS-containing ECS complexes, such as the ECS(SOCS7) complex, that regulate reelin signaling by mediating ubiquitination and degradation of DAB1 (By similarity). ECS(SOCS1) seems to direct ubiquitination of JAK2 (PubMed:11384984). The ECS(SOCS2) complex mediates the ubiquitination and subsequent proteasomal degradation of phosphorylated EPOR and GHR (PubMed:21980433, PubMed:25505247). The ECS(SPSB3) complex catalyzes ubiquitination of nuclear CGAS (PubMed:38418882). ECS(KLHDC1) complex is part of the DesCEND (destruction via C-end degrons) pathway and mediates ubiquitination and degradation of truncated SELENOS selenoprotein produced by failed UGA/Sec decoding, which ends with a glycine (PubMed:32200094). The ECS(ASB9) complex mediates ubiquitination and degradation of CKB (PubMed:33268465). As part of some ECS complex, promotes 'Lys-11'-linked ubiquitination and degradation of BTRC (PubMed:27910872). As part of a multisubunit ECS complex, polyubiquitinates monoubiquitinated POLR2A (PubMed:19920177). As part of the ECS(RAB40C) complex, mediates ANKRD28 ubiquitination and degradation, thereby inhibiting protein phosphatase 6 (PP6) complex activity and focal adhesion assembly during cell migration (PubMed:35512830). As part of the ECS(RAB40A) complex, mediates RHOU 'Lys-48'-linked ubiquitination and degradation, thus inhibiting focal adhesion disassembly during cell migration (PubMed:26598620). As part of the ECS(RAB40B) complex, mediates LIMA1/EPLIN and RAP2 ubiquitination, thereby regulating actin cytoskeleton dynamics and stress fiber formation during cell migration (PubMed:33999101, PubMed:35293963). May form a cell surface vasopressin receptor (PubMed:9037604).

Cellular Location

Nucleus. Note=Localizes to sites of DNA damage in a UBAP2 and UBAP2L-dependent manner.


KD-Validated Anti-Cullin 5 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

- [Flow Cytometry](#)
- [Cell Culture](#)

KD-Validated Anti-Cullin 5 Rabbit Monoclonal Antibody - Images



Western blotting analysis using anti-cullin 5 antibody (Cat#AGI2249). Total cell lysates (30 µg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-cullin 5 antibody (Cat#AGI2249, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.



Western blotting analysis using anti-cullin 5 antibody (Cat#AGI2249). Cullin 5 expression in wild-type (WT) and cullin 5 (CUL5) shRNA knockdown (KD) 293T cells with 20 µg of total cell lysates. Hsp90 α serves as a loading control. The blot was incubated with anti-cullin 5 antibody (Cat#AGI2249, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.